



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2021

---

## **A highly diverse bivalve fauna from a Bithynian (Anisian, Middle Triassic) Tubiphytes -microbial buildup in North Dobrogea (Romania)**

Friesenbichler, Evelyn ; Hautmann, Michael ; Grădinaru, Eugen ; Bucher, Hugo

**Abstract:** This paper describes a rich bivalve fauna from the Tubiphytes-Limestone Member (Bithynian, early middle Anisian) of the Caerace Formation in North Dobrogea (Romania). We report 51 bivalve species, including 7 new species and 2 new genera: *Atrina multicostata* sp. nov., *Pinna simionescui* sp. nov., *Chlamys* (*Praechlamys*) *prima* sp. nov., *Entolium reticulatum* sp. nov., *Scythentolium anisicum* sp. nov., *Romaniamya mahmudiaensis* gen. et sp. nov. and *Praedicerocardium vetulus* gen. et sp. nov. The latter taxon represents the oldest Triassic megalodontoid. We provide detailed descriptions of the bivalve fauna, discuss the ecology and compare species richness in the Early and Middle Triassic, showing that the bivalve assemblage of North Dobrogea is much more diverse than all previously described Early Triassic bivalve faunas, but not as rich as several faunas from the late Anisian and Ladinian. The relative richness of the Dobrogea fauna with respect to geologically older and younger faunas suggests that it represents an early stage of the main rediversification of bivalves after the end-Permian mass extinction.

DOI: <https://doi.org/10.1002/spp2.1286>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-175886>

Journal Article

Accepted Version

Originally published at:

Friesenbichler, Evelyn; Hautmann, Michael; Grădinaru, Eugen; Bucher, Hugo (2021). A highly diverse bivalve fauna from a Bithynian (Anisian, Middle Triassic) Tubiphytes -microbial buildup in North Dobrogea (Romania). *Papers in Palaeontology*, 7(1):447-495.

DOI: <https://doi.org/10.1002/spp2.1286>



**A highly diverse bivalve fauna from a Bithynian (Anisian, Middle Triassic) *Tubiphytes*-microbial buildup in North Dobrogea (Romania)**

Journal:	<i>Palaeontology</i>
Manuscript ID	PALA-10-18-4331-OA.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Friesenbichler, Evelyn; Universitat Zurich Palaontologisches Institut und Museum, Hautmann, Michael; University Zurich, Palaeontologisches Institut und Museum Grădinaru, Eugen; Universitatea din Bucuresti, Department of Geology Bucher, Hugo; Paläontologisches Institut und Museum, Universität Zürich;
Key words:	Bivalvia, Anisian, Bithynian, North Dobrogea, diversity, recovery

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**1 A highly diverse bivalve fauna from a Bithynian (Anisian, Middle Triassic) *Tubiphytes*-microbial**  
**2 buildup in North Dobrogea (Romania)**

4 Evelyn Friesenbichler<sup>1\*</sup>, Michael Hautmann<sup>1</sup>, Eugen Grădinaru<sup>2</sup>, Hugo Bucher<sup>1</sup>,

6 <sup>1</sup> Paläontologisches Institut und Museum, University of Zurich, Karl-Schmid-Strasse 4, 8006 Zurich,  
7 Switzerland; e-mails: evelyn.friesenbichler@pim.uzh.ch, michael.hautmann@pim.uzh.ch,  
8 hugo.fr.bucher@pim.uzh.ch

9 <sup>2</sup> Department of Geology, Faculty of Geology & Geophysics, University of Bucharest, Bd. Bălcescu  
10 Nicolae 1, RO-010041, Bucharest, Romania; e-mail: egradinaru@gmail.com

12 \*corresponding author

**15 Abstract**

17 This paper describes a rich bivalve fauna from the *Tubiphytes*-Limestone Member (Bithynian, early  
18 middle Anisian) of the Caerace Formation in North Dobrogea (Romania). We report 51 bivalve  
19 species, including seven new species and two new genera: *Atrina multicostata* n. sp., *Pinna*  
20 *simionescui* n. sp., *Chlamys (Praechlamys) prima* n. sp., *Entolium reticulatum* n. sp., *Scythentolium*  
21 *anisicum* n. sp., *Romaniamya mahmudiaensis* n. gen. et n. sp. and *Praedicerocardium vetulus* n. gen.

et n. sp. The latter taxon represents the oldest Triassic megalodontoid. We provide detailed descriptions of the bivalve fauna, discuss the ecology and compare species richness in the Early and Middle Triassic, showing that the bivalve assemblage of North Dobrogea is much more diverse than all previously described Early Triassic bivalve faunas, but not as rich as several faunas from the late Anisian and Ladinian. The relative richness of the Dobrogea fauna with respect to geologically older and younger faunas suggests that it represents an early stage of the main rediversification of bivalves after the end-Permian mass extinction.

**Keywords:** Bivalvia, Anisian, Bithynian, North Dobrogea, diversity, recovery.

The most severe biotic crisis in Earth's history, the end-Permian mass extinction about 252 Ma ago (Burgess *et al.* 2014), did not only extinguish between 81% (Stanley 2016) and 96% (Raup 1979) of all marine species, but also caused a temporal shift from the Upper Paleozoic T-type ('tropical') carbonate factory to a non-skeletal M-type ('mud-mound') carbonate factory (*sensu* Schlager 2000, 2003; Baud & Richoz 2004; Baud *et al.* 2007). While M-factory reefs already flourished during the Early Triassic (e.g. Brayard *et al.* 2011; Friesenbichler *et al.* 2018), T-factory reefs started re-diversifying since the Middle Triassic (Flügel & Stanley 1984; Stanley 1988; Senowbari-Daryan *et al.* 1993; Flügel 1994, 2002; Martindale *et al.* 2019). One of the last 'pure' M-type Triassic carbonate factories before the transition toward T-type factories is a *Tubiphytes*-microbial buildup located in the Mahmudia Quarry in the North Dobrogea Orogen (Popa *et al.* 2014), which is also one of the few bioconstructions from the Tethyan realm that flourished during the Bithynian (early middle Anisian, Middle Triassic).



1  
2  
3 46 The *in situ* boundstones of the *Tubiphytes*-microbial buildup developed on the upper slope of a distal  
4  
5 47 platform in relatively deep water under normal oxic conditions and in an intertropical environment  
6  
7 48 (Forel & Grădinaru 2018). It was built up by a micro-framework of *Tubiphytes*, a small encrusting and  
8  
9 49 low-tiering organism of uncertain affinities (Riding & Guo 1992) that is a common fossil in Anisian  
10  
11 50 reef assemblages. Apart from the micro-framework of *Tubiphytes*, the buildup contains abundant  
12  
13 51 fine-grained internal sediments, microbial crusts and autochthonous micrite, added by a large  
14  
15 52 volume of marine cements (Popa *et al.* 2014). The *Tubiphytes*-microbial buildup provided a suitable  
16  
17 53 environment for various groups of abundant and diverse benthic macroinvertebrates such as  
18  
19 54 bivalves, gastropods (Nützel *et al.* 2018) and brachiopods (Grădinaru & Gaetani 2019), plus a rich  
20  
21 55 ostracod fauna (Forel & Grădinaru 2018).  
22  
23  
24  
25  
26 56 The Bithynian *Tubiphytes*-microbial buildup is one of the oldest Triassic carbonate reefs and provides  
27  
28 57 the opportunity to study how bivalves adapted to this new habitat type and how this affected their  
29  
30 58 recovery after the end-Permian mass extinction. This study is part of a series of studies (Forel &  
31  
32 59 Grădinaru 2018; Nützel *et al.* 2018; Grădinaru & Gaetani 2019) that aim at the systematic description  
33  
34 60 and interpretation of the fossil content from the *Tubiphytes*-microbial buildup in North Dobrogea.  
35  
36 61 We report 51 bivalve species belonging to 39 genera, which makes bivalves the most diverse  
37  
38 62 macroinvertebrate group of the *Tubiphytes*-microbial buildup so far. Seven species and two genera  
39  
40 63 are newly described, including the oldest Triassic representative of the Megalodontoida. The ecology  
41  
42 64 of the bivalve fauna is reconstructed and its species richness is discussed in the context of the  
43  
44 65 rediversification of marine life after the end-Permian mass extinction.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

57 68 **GEOLOGICAL AND STRATIGRAPHICAL SETTING**  
58  
59 69  
60

70 The North Dobrogea Orogen is located in the foreland of the Alpine Carpathian Orogen (e.g.  
 71 Săndulescu 1995) and is limited by the Danube Delta in the North and the Black Sea in the East. It is  
 72 the westernmost segment of the Palaeo-Tethyan Cimmeride Orogenic System and continues  
 73 eastward into the Mountainous Crimea, the North Caucasus (Fig. 1A, inset map) and further  
 74 eastwards into the Asian Cimmerides (Săndulescu 1980; Şengör 1984, 1986). The North Dobrogea  
 75 Orogen is a fold-and-thrust belt that consists of several tectonic units (Fig. 1A. Visarion *et al.* 1990;  
 76 Săndulescu 1995). The Triassic succession has its widest extent in the Tulcea Unit, which is the  
 77 lowermost tectonic unit of the North Dobrogea Orogen and is located in its eastern part. The Triassic  
 78 sediments unconformably overlie the Variscan basement and are developed both in basinal and  
 79 carbonate platform facies (Grădinaru 1995, 2000). During the Middle Triassic, a carbonate platform  
 80 developed in the central and eastern areas of the Tulcea Unit. The various carbonate facies that  
 81 developed in the eastern area of the Tulcea Unit are described in detail by Popa *et al.* (2014). The  
 82 North Dobrogea Orogen, with its Tethyan-type Triassic succession, is tectonically bounded by the  
 83 Scythian Platform in the north and the Moesian Platform in the south (Fig. 1A), where the Triassic is  
 84 of Germanic type (Grădinaru 1995, 2000). The remote location of the North Dobrogea Tethyan-type  
 85 Triassic outside the Mediterranean Triassic is interpreted as a result of post-Triassic large-scale  
 86 horizontal displacements of Tethyan terranes due to the opening of the West Black Sea Basin (e.g.  
 87 Grădinaru 1988; Okay *et al.* 1994; Banks & Robinson 1997).

88 The studied material comes from the Mahmudia Quarry (45° 03' 12.41" N, 29° 03' 34.52" E), which is  
 89 located in the Caeracul Mare Hill area, south of the village of Mahmudia, 35 km east of Tulcea (Fig.  
 90 1A–B). The Middle Triassic Caerace Formation that is exposed in this limestone quarry can be  
 91 subdivided into the Stromatactis-Limestone Member and the *Tubiphytes*-Limestone Member (Fig. 1C  
 92 and Fig. 2). The latter is exposed in the northeastern part of the quarry and represents a  
 93 *Tubiphytes*-microbial buildup that was first documented by Dobre-Popa (2011) and Popa *et al.*  
 94 (2014). It yields a rich macroinvertebrate fauna consisting of coiled nautiloids, ammonoids, bivalves,  
 95 gastropods, brachiopods, sponges and crinoids (Popa *et al.* 2014; Forel & Grădinaru 2018; Nützel *et*

1  
2  
3 96 *al.* 2018; Grădinaru & Gaetani 2019). The ammonoids found in the *Tubiphytes*-Limestone Member  
4  
5 97 can be correlated with the Bithynian (early Middle Triassic) *Osmani*-ammonoid Zone of the  
6  
7 98 Northwestern Caucasus (Shevyrev 1995) and the lower part of the middle Anisian, i.e. the  
8  
9  
10 99 *Hyatti*-ammonoid Zone of northwestern Nevada (Silberling & Nichols 1982; Bucher 1992; Monnet &  
11  
12 100 Bucher 2006). The *Tubiphytes*-microbial buildup is a massive carbonate buildup without  
13  
14 101 well-expressed bedding and consists of several different carbonate facies (Popa *et al.* 2014). Among  
15  
16 102 these are frequent packages of microbially mediated cement crusts, which were generated by  
17  
18 103 syndepositional microbial mats. These microbial mats and their cement crusts provided taphonomic  
19  
20 104 conditions that were favourable for the preservation of minute details of the ornamentation of fossil  
21  
22 105 shells and the primary colour patterns of some gastropods (Popa *et al.* 2014; Forel & Grădinaru 2018;  
23  
24 106 Nützel *et al.* 2018). So far, only few data has been published on Triassic bivalves from the North  
25  
26 107 Dobrogea Orogen (Peters 1867; Kittl 1908; Simionescu 1908, 1910*a, b*, 1911, 1925; Mirăuță & Iordan  
27  
28 108 1982; Mirăuță *et al.* 1984), making the present work a significant contribution to the knowledge of  
29  
30 109 the bivalve fauna from this Tethyan-type Triassic development.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

35 110  
36  
37  
38 111  
39  
40  
41  
42 **MATERIAL AND METHODS**  
43  
44  
45  
46  
47

48 114 The bivalve fauna presented herein comes from one sample of c. 1.5 m<sup>3</sup> rock material from the  
49  
50 115 *Tubiphytes*-Limestone Member (Fig. 1C–D and Fig. 2) that yielded several hundreds of  
51  
52 116 macroinvertebrate fossils. For taxonomical identification, a total of 303 bivalve specimens were  
53  
54 117 mechanically prepared from the rock with micro-jacks. The terms small, medium sized and large are  
55  
56 118 used for specimens that are smaller than 2 cm, between 2 and 5 cm or larger than 5 cm, respectively.  
57  
58  
59 119 The 303 specimens represent a minimum of 214 individuals. This number was inferred from counting  
60

the most numerous valves, left valves (LVs) or right valves (RVs), together with articulated specimens. The number of individuals was the basis for a rarefaction analysis (Sanders 1968), which was performed to assess the relative completeness of sampling and to compare the diversities of the present bivalve fauna with other Early and Middle Triassic bivalve assemblages. Ecological parameters like Simpson's index of dominance (Simpson 1949) and rarefaction analysis were calculated with PAST version 2.17 software (Hammer *et al.* 2001). For further taxonomical as well as ecological characterization of the bivalve fauna, rank-abundance distributions (Whittaker 1965) and pie charts were used. The modes of life were inferred from literature data, functional morphology and the mode of life of related Recent taxa. For photographing, specimens were covered with ammonium chloride. The comparison of diversities of Early and Middle Triassic bivalve faunas is based a previous analysis (Friesenbichler *et al.* 2019) and additional literature. The occurrences of identified bivalve species and genera during the Early and Middle Triassic was analysed based on the literature including the compendium of Ros-Franch *et al.* (2014). The bivalve samples are housed in the Collection of the Museum of the Laboratory of Palaeontology in the University of Bucharest.

## SYSTEMATIC PALAEONTOLOGY

Our systematic arrangement is based on Cox *et al.* (1969), Hautmann (2001a) and Bieler *et al.* (2014), with additions from Allasinaz (1972), Waller & Stanley (2005), Hautmann (2008), Carter *et al.* (2011), Hautmann *et al.* (2013) and Hofmann *et al.* (2014).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

142  
  
  
  
143  
  
  
  
144  
  
  
  
145  
  
  
  
146  
  
  
  
147  
  
  
  
148  
  
  
  
149  
  
  
  
  
  
  
  
150  
  
  
  
  
  
  
  
151  
  
  
  
152  
  
  
  
  
  
  
  
153  
  
  
  
154  
  
  
  
155  
  
  
  
156  
  
  
  
157  
  
  
  
158  
  
  
  
159  
  
  
  
160  
  
  
  
161

Class BIVALVIA Linnaeus, 1758

Subclass PTERIOMORPHIA Beurlen, 1944

Order ARCOIDA Stoliczka, 1871 in Stoliczka 1870–1871

Superfamily ARCOIDEA de Lamarck, 1809

Family PARALLELODONTIDAE Dall, 1898

Genus PARALLELODON Meek & Worthen, 1866

*Type species. Macrodon rugosus* Buckman in Murchison *et al.*, 1845.

*Parallelodon?* sp.

Figures 3A–B

*Material.* One incompletely preserved LV.

*Description.* Shell medium sized, posteriorly elongated, moderately inflated; dorsal margin long and straight, posterior margin slightly concave, ventral margin convex; ligament duplivincular, posterior part of ligament area with subparallel grooves representing traces of duplivincular ligament system (Fig. 3B); carina rounded, extending to postero-ventral part of shell, bordering slightly impressed posterior area; irregular growth lamellae most clearly visible near shell margins.

*Remarks.* The shape, outline and duplivincular ligament area indicate that this specimen belongs to the Arcoida. Among Middle Triassic arcoids, three species are similar in terms of the elongated outline and the presence of an irregular concentric sculpture: *Parallelodon beyrichi* (von Strombeck, 1849), *P. esinensis* (Stoppani, 1858) and *P. impressum* (Münster, 1841). In *P. esinensis* this concentric sculpture is best visible at the shell margins. Furthermore, it possesses a rounded carina, which is also the case in the Dobrogea specimen. However, Posenato (2008) mentions a very shallow but wide sulcus in the middle part of the shell in front of the carina, which is not observable in the Dobrogea specimen because of its incomplete preservation. Apart from that, the antero-ventral margin of *P. esinensis* is more rounded compared to the antero-ventral margin of the Dobrogea specimen, which is more similar to the antero-ventral margin of *P. beyrichi*. However, like *P. esinensis*, *P. beyrichi* possesses a shallow median sulcus that runs from the beak to the ventral margin. Furthermore, radial ribs were reported on the area of *P. beyrichi*, a feature that is not observable in the Dobrogea specimen. As well as *P. beyrichi*, *P. impressum* possesses a sulcus and radial ribs. However, this sulcus is only present in juvenile specimens and the radial ribs are neither observable with naked eye nor are they present in all specimens of *P. impressa*. Already Salomon (1895, p. 163–165) pointed out that *P. impressum* and *P. beyrichi* cannot be clearly distinguished from each other; therefore he united both species as *P. impressum*. Furthermore, he proposed that *P. impressum* could possibly represent juvenile specimens of *P. esinensis*. Currently, the problem of possible synonymies cannot be resolved and an assignment of the Dobrogea specimen to any of these species is uncertain. Therefore we present it in open nomenclature and provisionally assign it to the genus *Parallelodon*.

*Ecology.* According to Stanley (1972), the genus *Parallelodon* comprises endobenthic as well as epibenthic species. Yin & McRoberts (2006) suggested that species of this genus had an epibyssate and suspensivorous mode of life.

1		
2		
3	187	
4		
5		
6	188	Order MYTILIDA Férussac, 1822 in Férussac 1821–1822
7		
8		
9	189	Superfamily MYTILOIDEA Rafinesque, 1815
10		
11		
12	190	Family MYTILIDAE Rafinesque, 1815
13		
14		
15	191	Genus MODIOLUS de Lamarck, 1799
16		
17	192	
18		
19		
20	193	<i>Type species. Mytilus modiolus</i> Linnaeus, 1758.
21		
22		
23		
24	194	
25		
26		
27	195	<i>Modiolus</i> sp. A
28		
29	196	Figures 3C–D
30		
31		
32	197	
33		
34		
35	198	<i>Material.</i> Two incompletely preserved LVs.
36		
37		
38	199	
39		
40		
41	200	<i>Description.</i> Shell medium sized, modioliform, retrocrescent, inequilateral; dorsal margin slightly
42		
43	201	convex, passing gradually into convex posterior margin, ventral margin concave; beak acute,
44		
45	202	prosogyrate, subterminal; byssal gape well developed (Fig. 3D), located at one third of antero-ventral
46		
47	203	margin; anterior shell lobus small; irregular growth lamellae, best visible at posterior and ventral
48		
49	204	margin.
50		
51		
52		
53	205	
54		
55		
56	206	<i>Remarks.</i> The shape, outline and the presence of a byssal gape are indicative for the genus <i>Modiolus</i> .
57		
58	207	Already Bittner (1901, p. 21) noted the difficulties in differentiating between species among Triassic
59		
60		

208 representatives of this morphologically variable genus. Because of its incomplete preservation, we  
209 present the Dobrogea specimens in open nomenclature.

210

211 *Ecology.* Endobyssate suspension-feeder (Stanley 1970, 1972).

212

213 *Modiolus* sp. B

214 Figure 3E

215

216 *Material.* One incompletely preserved LV.

217

218 *Description.* Shell small, modioliform, slightly inflated; beak acute, prosogyrate, subterminal;  
219 antero-ventral margin nearly straight, posterior margin curved; very fine and regular growth lines.

220

221 *Remarks.* This taxon differs from *Modiolus* sp. A in being less upright and having a straight  
222 antero-ventral margin and fine, regular growth lines.

223

224 *Ecology.* Same as for *Modiolus* sp. A.

225



226 Order PTERIOIDA Newell, 1965

227 Suborder PTERIINA Newell, 1965

228 Superfamily PINNOIDEA Leach, 1819

229 Family PINNIDAE Leach, 1819

230

231 Remarks: Three genera of Pinnidae have been reported from the Triassic: *Pinna* Linnaeus, 1758,  
232 *Atrina* Gray, 1842 and *Mesopinna* Vu Khuc & Huyen, 1984. One character used for the distinction of  
233 *Pinna* and *Atrina* is the shape of the nacreous region on the shell interior. The nacre in *Pinna* is  
234 medially divided into two lobes, whereas the nacreous region in *Atrina* is continuous (Cox *et al.*  
235 1969). Another commonly used criterion for distinguishing *Pinna* and *Atrina* is the presence or  
236 absence of a medial carina. Accordingly, *Atrina* does not possess a medial carina, which is present in  
237 *Pinna*, at least in earlier growth stages (Cox *et al.* 1969; Waller & Stanley 2005; Koppka 2018).  
238 However, according to Hyatt (1892) and Turner & Rosewater (1958) both genera show little  
239 morphological difference with respect to the medial carina during their early ontogenetic stages.  
240 Therefore, the differentiation of both genera based on the presence or absence of a medial carina is  
241 only possible for adult specimens. Moreover, only few Triassic species assigned to *Pinna*  
242 demonstrably possess a medial carina (Waller & Stanley 2005; McRoberts 2017). This could mean  
243 either that these species do not belong to *Pinna*, or that the presence of a medial carina does not  
244 sufficiently diagnose this genus. Another character used to distinguish *Pinna* from *Atrina* is the shape.  
245 According to Cox *et al.* (1969) and Waller & Stanley (2005), *Atrina* is ham-shaped with a rounded  
246 posterior end, whereas *Pinna* is wedge- to ham-shaped. *Mesopinna*, the third Triassic pinniid genus,  
247 was described by Vu Khuc (1991) as pinniform (i.e. wedge-shaped), with a weakly pronounced medial  
248 carina, a lenticular shell cross section at the umbonal region and a concentric sculpture.

249 The material described below demonstrates the difficulty of a consistent separation of *Pinna* and  
250 *Atrina* based solely on one criterion. Since the nacreous regions of the shell's interior are not

observable, we cannot use this criterion for the differentiation between *Pinna* and *Atrina*, but we found that a more pronounced carina correlates with a more wedge-shaped outline, whereas a short carina corresponds to a ham-shaped overall morphology. We therefore suggest tentatively that *Pinna* is wedge-shaped to slightly ham-shaped and has a medial carina that was present during most of its ontogeny, whereas *Atrina* is clearly ham-shaped, with a medial carina either absent or present only during its earliest ontogeny. The main difference between *Pinna* and *Mesopinna* is the absence of radial ribs in *Mesopinna*, which, in our opinion, is insufficient to justify its separation from *Pinna*.

#### Genus ATRINA Gray, 1842

*Type species. Pinna nigra* Dillwyn, 1817.

#### *Atrina multicostata* n. sp.

Figure 3H

*LSID.* urn:lsid:zoobank.org:act:41925AAC-F737-41D4-8631-2AB10A18BCF7

*Derivation of name.* Combination of *multi* (Latin: many) and *costata* (Latin: ribbed).

*Holotype.* LPB IIL 2005

*Material.* One LV.

1  
2  
3 273  
4  
5  
6 274 *Diagnosis.* *Atrina* with at least 40 radial ribs with rounded crests that get broader towards posterior  
7  
8 275 margin.  
9  
10  
11 276  
12  
13  
14 277 *Description.* Shell large, inequilateral, ham-shaped, weakly inflated; dorsal margin long and straight,  
15  
16 278 antero-ventral margin concave, posterior margin regularly rounded; beak obtuse and terminal; carina  
17  
18 279 well defined, present on anterior part of shell, fading out distally; at least 40 radial ribs with rounded  
19  
20 280 crests, regularly intercalated, getting broader towards posterior margin, straight on dorsal part of  
21  
22 281 valve but curved on ventral part, interspaces as broad as or slightly broader than ribs, getting  
23  
24 282 broader towards posterior margin; internal structures unknown.  
25  
26  
27  
28  
29 283  
30  
31  
32 284 *Remarks.* Because of the ham-shaped outline and a medial carina that was only present during early  
33  
34 285 ontogeny, we assign this specimen to *Atrina* (see above). So far, *Atrina* was reported from the  
35  
36 286 Triassic only by Waller & Stanley (2005) and Stiller (2001). In contrast to the Dobrogea species, *Atrina*  
37  
38 287 *sinuata* Waller & Stanley, 2005 possesses commarginal ridges. Stiller (2001; p. 277) mentions *Atrina*  
39  
40 288 n. sp. A from the Leidapo Member (late Pelsonian – early Illyrian, China), without providing a  
41  
42 289 description, but judging from the described material that he assigns to this species, it lacks the  
43  
44 290 numerous radial ribs that characterize *Atrina multicostata*.  
45  
46  
47  
48 291 The sculpture of *Atrina multicostata* is reminiscent of the Jurassic *Stegoconcha* Böhm, 1907, a genus  
49  
50 292 that is closely related to *Atrina* but differs, among other things, in being higher than long.  
51  
52  
53  
54 293  
55  
56  
57 294 *Ecology.* *Atrina* is a semi-infaunal suspension-feeder that is byssally attached to coarse sediment  
58  
59 295 particles. However, in contrast to *Pinna* that lives with two-thirds of its shell into the sediment, *Atrina*  
60

296 lives almost completely buried (Garcia-March *et al.* 2008). This mode of live is also assumed for  
297 Mesozoic species of that genus (Ros-Franch *et al.* 2014).

298

299 Genus PINNA Linnaeus, 1758

300

301 *Type species. Pinna rudis* Linnaeus, 1758.

302

303 *Pinna simionescui* n. sp.

304 Figure 3G

305

306 *LSID.* urn:lsid:zoobank.org:act:1A562F69-FB98-45FD-98F4-4DF4B6E1FE9E

307

308 *Derivation of name. Simionescui*, after the Romanian palaeontologist Ion Simionescu, for his  
309 contribution to the Triassic of North Dobrogea.

310

311 *Holotype.* LPB IIL 2006

312

313 *Material.* One RV.

314

315 *Diagnosis.* *Pinna* with regularly arranged commarginal shell folds.

1  
2  
3 316  
4  
5  
6 317 *Description.* Shell medium sized, inequilateral, subtrigonal, retrocrescent, moderately inflated at the  
7  
8 318 anterior part of the valve and becoming flatter towards posterior margin; dorsal margin straight,  
9  
10 319 ventral margin concave; beak terminal and acute; carina well developed in anterior part of the shell  
11  
12  
13 320 but fading away in the posterior part, separating valve into two parts of approximately equal size;  
14  
15 321 regular commarginal shell folds; internal structures unknown.  
16  
17  
18 322  
19  
20  
21 323 *Remarks.* Because of the wedge- to slightly ham-shaped outline and a medial carina that was present  
22  
23 324 during most of the ontogeny we assign this species to *Pinna* (see above). *Pinna yunnanensis* Chen,  
24  
25 325 1976 from the Middle and Late Triassic of China is similar in the presence of commarginal folds;  
26  
27 326 however, these are broader and stronger developed compared to the *Dobrogea* specimen.  
28  
29 327 Moreover, *P. yunnanensis* differs from the *Dobrogea* specimen by its straight ventral margin,  
30  
31 328 resulting in a cuneiform shape, and by a continuously developed carina. *Pinna mileensis* Guo, 1985 is  
32  
33 329 another pinnid with commarginal ribs that was reported from the Middle Triassic of China. In  
34  
35 330 contrast to *Pinna simionescui*, the commarginal ribs of *P. mileensis* are finer but higher. Furthermore,  
36  
37 331 the ventral margin is straight and the carina is continuously developed. All other Triassic species of  
38  
39 332 this genus, e.g. *P. ecki* Picard, 1907, *P. meriani* Winkler, 1859, *P. tommasii* von Wöhrmann & Koken,  
40  
41 333 1892 and *P. vomis* Winkler, 1859, differ significantly in their sculpture from the *Dobrogea* specimen,  
42  
43 334 which we therefore describe as a new species.  
44  
45  
46  
47  
48  
49 335  
50  
51  
52 336 *Ecology.* Species of the genus *Pinna* are semi-infaunal suspension-feeders that are byssally attached  
53  
54 337 to larger sediment particles (Cox *et al.* 1969; McRoberts 2017).  
55  
56  
57 338  
58  
59  
60

339 *Pinna?* cf. *raibliana* Parona, 1889

340 Figure 3I

341

342 cf. \*1889 *Pinna raibliana*; Parona, p. 105, pl. 8, fig. 11.

343 cf. 1923 *Pinna raibliana* Parona; Diener, p. 87 [cum syn.].

344 cf. 1966 *Pinna (Pinna) raibliana* Parona; Allasinaz, p. 649, pl. 41, fig. 15.

345 cf. 1979 *Pinna raibliana* Parona; Lieberman, p. 100, pl. 4, fig. 8.

346

347 *Material.* One fragmented internal mould, probably from a RV.

348

349 *Description.* Shell medium sized, elongated, cuneiform and relatively flat; at least six radial ribs.

350

351 *Remarks.* Based on the ornamentation, the Dobrogea specimen could belong to *Pinna raibliana* that

352 possesses 13 to 15 radial and nodose ribs on its dorsal parts. *P. raibliana* was described as strongly

353 inflated (Parona 1889), which cannot be verified in the incomplete specimen from Dobrogea.

354 Although the radial ribs indicate a wedge-shaped outline, the incomplete preservation prevents us

355 from confirming the presence or absence of a medial carina. Our taxonomic assignment is therefore

356 tentative. *Pinna?* cf. *raibliana* differs from *Pinna simionescui* by the presence of radial sculptural

357 elements and the absence of commarginal shell folds.

358

359 *Ecology.* Same as for *Pinna simionescui*.

1		
2		
3	360	
4		
5		
6	361	<i>Occurrence.</i> Acquate (Italy, Julian; Parona 1889), Sella delle Cave (Italy, Julian; Allasinaz 1966),
7		
8	362	Conzen Formation (Italy, Julian; Lieberman 1979), Valletta di Rogno (Italy, Carnian; Allasinaz 1972).
9		
10		
11	363	
12		
13		
14		
15	364	Superfamily POSIDONIOIDEA Neumayr, 1891
16		
17		
18	365	Family POSIDONIIDAE Frech, 1909
19		
20		
21	366	Genus BOSITRA de Gregorio, 1886
22		
23	367	
24		
25		
26	368	<i>Type species.</i> <i>Posidonia ornati</i> Quenstedt, 1851 in Quenstedt 1851–1852.
27		
28		
29	369	
30		
31		
32	370	<i>Bositra wengensis</i> (Wissmann, 1841)
33		
34	371	Figure 3F
35		
36		
37	372	
38		
39		
40		
41	373	*1841 <i>Posidonomya wengensis</i> ; Wissmann, p. 23, pl. 16, fig. 12.
42		
43		
44	374	1858 <i>Posidonomya wengensis</i> Wissmann; Stoppani, p. 95, pl. 19, fig. 12.
45		
46		
47	375	1865 <i>Posidonomya wengensis</i> Wissmann; Laube, p. 76, pl. 20, fig. 12.
48		
49		
50	376	1889 <i>Posidonomya wengensis</i> Wissmann; Parona, p. 106, pl. 8, fig. 10.
51		
52		
53	377	1904 <i>Posidonia wengensis</i> mut. nov. <i>alta</i> (Wissmann); Frech, p. 16, fig. 15.
54		
55		
56	378	1923 <i>Posidonia wengensis</i> (Wissmann); Diener, p. 64 [cum syn.].
57		
58		
59	379	1976 <i>Posidonia wengensis</i> (Wissmann); Gu <i>et al.</i> , pl. 34, figs 32–34.
60		

1981 *Posidonomya wengensis* Wissmann; Zardini , pl. 21, figs 7–8.

2003 *Posidonia wengensis* (Wissmann); Szente & Vörös, p. 127, pl. Biv-I, figs 27–28 [cum syn.].

382

383 *Material.* One RV.

384

385 *Description.* Shell small, thin, ovate, longer than high, weakly inflated, infracrescent, nearly  
 386 equilateral; beak orthogyrate, obtuse, not projecting above straight dorsal margin; dorsal margin  
 387 without auricles, antero-dorsal margin slightly shorter than posterior one; regular commarginal ribs.

388

389 *Remarks.* ‘*Daonella*’ *boeckhi* Mojsisovics, 1874 is morphologically similar, but possesses a radial  
 390 ornamentation in addition to commarginal ribs. The outline, presence of commarginal ribs, absence  
 391 of a radial ornamentation and absence of auricles are diagnostic features of *Bositra*.

392

393 *Ecology.* Several modes of life have been suggested for *Bositra*, such as pseudoplanktonic (Stanley  
 394 1972), nekto planktonic (Jefferies & Minton 1965; Hayami 1969; Duff 1975) or benthic (Conti &  
 395 Monari 1992; Etter 1996). Etter (1996) showed that a benthic mode of life is most likely, and  
 396 provided arguments against a pseudoplanktonic or nekto planktonic mode of life. Since no byssal  
 397 notch has been observed, Waller & Stanley (2005) suggested a reclined mode of life for *Bositra*.

398

399 *Occurrence.* Leidapo Member (China, late Pelsonian – early Illyrian; Komatsu *et al.* 2004), Felsőőrs  
 400 Formation (Hungary, Pelsonian–Illyrian; Szente & Vörös 2003), Nam Tham Formation (Vietnam, early  
 401 Ladinian; Vu Khuc 2000), Cassian Formation (Italy, Longobardian–Julian; Laube 1865; Bittner 1895;



1  
2  
3 402 Zardini 1981), Esino Limestone (Italy, Ladinian; Stoppani 1858), Muong Trai Formation (Vietnam,  
4  
5 403 Ladinian; Vu Khuc 2000), Hon Nghe Formation (Vietnam, Ladinian; Vu Khuc 2000), Hong Hoi  
6  
7 404 Formation (Thailand, Ladinian; Vu Khuc 2000), Jiangdonggou Formation (China, Middle Triassic; Gu *et*  
8  
9  
10 405 *al.* 1976), Cataloi Formation (Romania, Ladinian–Carnian; Turculet 2009), Acquate (Italy, Julian;  
11  
12 406 Parona 1889).

13  
14  
15 407

16  
17  
18 408 Superfamily PTERIOIDEA Gray, 1847

19  
20  
21 409 Family PTERIIDAE Gray, 1847

22  
23  
24 410 Genus PTERIA Scopoli, 1777

25  
26  
27 411

28  
29  
30 412 *Type species. Mytilus hirundo* Linnaeus, 1758.

31  
32  
33 413

34  
35  
36 414 *Remarks.* The general morphology of the following species is indicative for the Pteriidae as well as  
37  
38 415 the Bakevelliidae. A definite assignment to one of these families is not possible because the internal  
39  
40 416 shell morphology including hinge and ligament is unknown. We assign it to *Pteria* because this  
41  
42 417 corresponds to its original assignment to ‘*Avicula*’ (= *Pteria*).

43  
44  
45 418

46  
47  
48 419 *Pteria sturi* (Bittner, 1895)

49  
50  
51 420 Figures 3J–K

52  
53  
54 421

55  
56  
57 422 1865 *Avicula gea* d’Orbigny; Laube, p. 50, pl. 16, fig. 9.

58  
59  
60 423 1889 *Avicula gea* d’Orbigny; Parona, p. 93, pl. 8, fig. 7.

424 \*1895 *Avicula sturi*; Bittner, p. 69, pl. 8, figs 1–4.

425 1923 *Avicula sturi* Bittner; Diener, p. 25 [cum syn.].

426 1966 *Pteria sturi* (Bittner); Allasinaz, p. 652, pl. 42, figs 4–8 [cum syn.].

427 1979 *Pteria sturi* (Bittner); Lieberman, p. 96, pl. 3, fig. 5 [cum syn.].

428 1981 *Arcavicula sturi* (Bittner); Zardini, pl. 8, fig. 15–16; pl. 10, figs 8–9.

429 2004 *Pteria cf. sturi* (Bittner); Komatsu *et al.*, fig. 7(5).

430 2010 *Pteria sturi* (Bittner); Komatsu *et al.*, figs 4(2–4).

431

432 *Material.* Four LVs.

433

434 *Description.* Shell medium sized, pteriiform, longer than high, oblique, weakly inflated; beak of LV  
 435 acute, prosogyrate, slightly projecting above straight and long dorsal margin; anterior auricle small  
 436 and not well demarcated from flank; shell margin folded inward below anterior auricle, forming a  
 437 byssal gape; posterior wing long, posteriorly pointed, demarcated from flank by deep furrow;  
 438 elongated groove on posterior wing running subparallel to the dorsal margin; growth lines of variable  
 439 strength.

440

441 *Remarks.* As noted by Bittner (1895, p. 69), the specimens described as *Avicula gea* d’Orbigny, 1849  
 442 by Laube (1865) and subsequent authors belong to a different species, for which he introduced the  
 443 new name *Avicula sturi*. The features of the Dobrogea specimen match those of *Avicula sturi* as  
 444 described by Bittner (1895) and Allasinaz (1966), particularly with regard to shell shape, pointed  
 445 posterior wing and presence of a furrow that extends below the postero-dorsal margin.

1  
2  
3 446  
4  
5  
6 447 *Ecology*. Recent species of the genus *Pteria* live epibyssate, preferentially attached to corals (Stanley  
7  
8 448 1970, 1972) and plants (Cox *et al.* 1969, figs. 87–88). However, no corals or plants are reported from  
9  
10 449 the *Tubiphytes*-microbial buildup, thus it is uncertain whether the Dobrogea specimens were  
11  
12  
13 450 attached to such organisms or to some other kind of hard substratum.  
14  
15  
16 451  
17  
18  
19 452 *Occurrence*. Leidapo Member (China, late Pelsonian – early Illyrian; Komatsu *et al.* 2004), Na Khuat  
20  
21 453 Formation (North Vietnam, Anisian–Ladinian; Komatsu *et al.* 2010), Cassian Formation (Italy,  
22  
23 454 Longobardian–Julian; Bittner 1895; Fürsich & Wendt 1977; Zardini 1981), Sella delle Cave (Italy,  
24  
25 455 Julian; Allasinaz 1966), Rio del Lago Formation (Italy, Upper Julian; Lieberman 1979), Tor Formation  
26  
27 456 (Italy, Upper Julian–Tuvanian; Lieberman 1979).  
28  
29  
30  
31 457  
32  
33  
34 458 *Pteria cf. cassiana* (Bittner, 1895)  
35  
36 459 Figure 3N  
37  
38  
39 460  
40  
41  
42 461 *cf.* \*1895 *Avicula cassiana*; Bittner, p. 71, pl. 8, figs 6–8.  
43  
44  
45 462 *cf.* 1923 *Avicula cassiana* Bittner; Diener, p. 18 [cum syn.].  
46  
47  
48 463 *cf.* 1976 *Pteria cassiana* (Bittner); Qihong *et al.*, pl. 27, figs 40–42, 45–46.  
49  
50  
51 464 *cf.* 1981 *Arcavicula cassiana* (Bittner); Zardini, pl. 10, figs 11a–b.  
52  
53  
54 465  
55  
56  
57 466 *Material*. One incompletely preserved interior mould of a RV with some shell remains.  
58  
59  
60 467

468 *Description.* Shell medium sized, pteriiform, approximately as long as high, weakly inflated; dorsal  
469 margin straight and long; anterior auricle not preserved, posterior auricle elongated, acute, with  
470 broad and shallow sinus; more or less regular growth lines.

471

472 *Remarks.* The outline of this taxon is almost identical to that of *Avicula cassiana* Bittner, 1895.  
473 However, the presence of up to three commarginal ribs on its posterior auricle that characterizes *A.*  
474 *cassiana* cannot be verified in the Dobrogea specimen. The difference between this species and  
475 *Pteria sturi* is the broader and shallower sinus below the posterior auricle in *P. sturi*. Furthermore, *P.*  
476 *cf. cassiana* is more upright than *P. sturi*.

477

478 *Ecology.* Same as for *Pteria sturi*.

479

480 *Occurrence.* Leidapo Member (China, late Pelsonian – early Illyrian; Stiller 2001; Komatsu *et al.* 2004),  
481 Cassian Formation (Italy, Longobardian–Julian; Bittner 1895; Fürsich & Wendt 1977; Zardini 1981),  
482 Weiyuanjiang Formation (China, Carnian; Qihong *et al.* 1976).

483

484 *Pteria* sp. A

485 Figure 3L

486

487 *Material.* One fragmented LV and one incompletely and strongly recrystallized RV.

488

1  
2  
3 489 *Description.* Shell medium sized, pteriiform, longer than high, weakly inflated, inflation strongest at  
4  
5 490 umbonal region and flat at postero-ventral region; beak acute, prosogyrate, slightly projecting above  
6  
7 491 straight and long dorsal margin; anterior auricle not observed, posterior auricle larger, triangular, not  
8  
9 492 pointed.  
10  
11  
12  
13 493  
14  
15  
16 494 *Remarks.* The incomplete preservation of the specimen prevents a determination at the species level.  
17  
18 495 The specimen is similar to *Pteria sturi* but the posterior wing lacks the pointed distal end that  
19  
20 496 characterizes this species.  
21  
22  
23 497  
24  
25  
26 498 *Ecology.* Same as for all other *Pteria* species.  
27  
28  
29 499  
30  
31  
32  
33 500 *Pteria* sp. B  
34  
35 501 Figure 3M  
36  
37  
38 502  
39  
40  
41 503 *Material.* One partly preserved RV.  
42  
43  
44 504  
45  
46  
47 505 *Description.* Shell medium sized, pteriiform, strongly inflated; beak terminal, acute, prosogyrate,  
48  
49 506 slightly projecting above straight dorsal margin; posterior auricle well demarcated from body shell;  
50  
51 507 radial lines and growth lines.  
52  
53  
54  
55 508  
56  
57  
58 509 *Remarks.* The outline and general features of this specimen are reminiscent of *Gervilleia? ogilviae*  
59  
60 510 Bittner, 1895, including the presence of radial lines and growth lines, but the growth lines of *G.?*

511 *ogilviae* are stronger and not as numerous as in the Dobrogea specimen. Bakevellidae indet. reported  
512 from Foster *et al.* (2019; Fig. 7F) also possesses radial ribs, but this taxon is represented by an  
513 incompletely preserved specimen and does not allow a detailed comparison. *Pteria* sp. B differs from  
514 the other Dobrogea *Pteria* species by the radial sculptural elements and the stronger inflation.

516 *Ecology.* Same as for all other *Pteria* species.

518 Family BAKEVELLIIDAE King, 1850

519 Genus CULTRIOPSIS Cossmann, 1904

521 *Type species.* *Gervillia (Cultriopsis) falciformis* Cossmann, 1904.

523 *Cultriopsis* sp.

524 Figure 4A

526 *Material.* One incompletely preserved LV.

528 *Description.* Shell medium sized, lanceolate, retrocrescent, inequilateral; beak acute, prosogyrate,  
529 terminal; irregular growth lamellae on ventral margin; anterior part incompletely preserved.

530

1  
2  
3 531 *Remarks.* In spite of its incomplete preservation, this specimen shows the distinctive shape of  
4  
5 532 *Cultriopsis*. However, a determination at the species level is impossible because the posterior auricle  
6  
7 533 is not preserved.  
8  
9  
10  
11 534  
12  
13  
14 535 *Ecology.* Byssally attached epifaunal suspension-feeder (Hautmann 2001*a*).  
15  
16  
17 536  
18  
19  
20 537 Genus GERVILLARIA Cox, 1954  
21  
22 538  
23  
24  
25 539 *Type species.* *Modiola? alaeformis* J. Sowerby, 1819.  
26  
27  
28 540  
29  
30  
31 541 *Gervillaria subelegans* (Chen in Gu *et al.*, 1976)  
32  
33 542 Figures 4B–C  
34  
35  
36 543  
37  
38  
39 544 \*1976 *Bakevellidoes subelegans*; Chen, p. 136, pl. 27, figs 8–9.  
40  
41  
42 545 1976 *Bakevellidoes subelegans* Chen; Qihong *et al.*, pl. 28, fig. 38.  
43  
44  
45 546 2004 *Gervillaria subelegans* (Chen); Komatsu *et al.*, p. 216, fig. 7(4).  
46  
47  
48 547  
49  
50  
51 548 *Material.* Three LVs.  
52  
53  
54  
55 549  
56  
57  
58 550 *Description.* LV medium to large in size, higher than long, retrocrescent, strongly inflated, bialate;  
59  
60 551 beak prosogyrate, acute, rising prominently above straight dorsal margin; anterior auricle weakly

552 demarcated from main body of shell, posterior auricle larger than anterior one, better demarcated  
553 from main body of shell by sulcus and steep diagonal carina; regular growth lines on the entire  
554 surface of shell.

555

556 *Remarks.* Although the internal characters of the Dobrogea specimens are unknown, they agree very  
557 well with *Gervillaria subelegans*, as illustrated in Komatsu *et al.* (2004), in terms of size, overall  
558 morphology and ornamentation.

559

560 *Ecology.* *Gervillaria* has been interpreted as an epifaunal or semi-infaunal recliner (Muster 1995, p.  
561 99; Aberhan & Muster 1997).

562

563 *Occurrence.* Leidapo Member (China, late Pelsonian – early Illyrian; Stiller 2001; Komatsu *et al.* 2004),  
564 Yunnan (China, Middle Triassic; Qihong *et al.* 1976).

565

566 Genus UNCERTAIN

567

568 Bakevelliidae indet.

569 Figure 4D

570

571 *Material.* One incompletely preserved LV.

572



1  
2  
3 573 *Description.* Shell large, pteriiform, longer than high, strongly inflated; beak prosogyrate, projecting  
4  
5 574 above straight dorsal margin; anterior auricle not preserved, posterior auricle well demarcated from  
6  
7 575 flank.  
8  
9  
10  
11 576  
12  
13  
14 577 *Remarks.* The shape and the strong inflation indicate that this specimen belongs to the Bakevelliidae,  
15  
16 578 but due to incomplete preservation a more accurate determination is not possible.  
17  
18  
19 579  
20  
21  
22 580 *Ecology.* Probably the same as for *Gervillaria subelegans*.  
23  
24  
25 581  
26  
27  
28 582 Suborder PECTININA Waller, 1978  
29  
30  
31 583 Superfamily AVICULOPECTINOIDEA Meek & Hayden, 1865  
32  
33  
34 584 Family UNCERTAIN  
35  
36  
37 585  
38  
39  
40 586 *Remarks.* The two taxa described below have a pectinoid shape but lack a ctenolium, which is a key  
41  
42 587 character of Pectinoidea in the strict sense (Waller 1984; Hautmann 2010). We therefore assign  
43  
44 588 these taxa to the Aviculopectinoidea s. l., which corresponds to the hyporder Aviculopectinoidei  
45  
46 589 excluding the Pterinopectinoidea in the classification of Carter *et al.* (2011). Although  
47  
48 590 Aviculopectinoidei is heavily oversplit, the combination of characters observed in these taxa does not  
49  
50  
51 591 conform to the character set of any available genus. This problem raises the general question of  
52  
53 592 classification in a clade that is characterized by extreme morphological convergence, plasticity, and  
54  
55 593 mosaic evolution, which is beyond the scope of the present paper. We therefore present these taxa  
56  
57 594 in open nomenclature and will discuss their taxonomy in a future study.  
58  
59  
60 595

596 'Pecten' *volaris* (Bittner, 1902)

597 Figure 4H

598

599 \*1902 *Pecten volaris*; Bittner, p. 634, pl. 27(10), fig. 32.

600 1972 *Chlamys* (*Chlamys*) *volaris* (Bittner); Allasinaz, p. 339, pl. 44, fig. 9; pl. 45, figs 1–2.

601 1981 cf. *Chlamys volaris* (Bittner); Zardini, pl. 18, figs 8a–b.

602

603 *Material*. One LV and one internal mould of a LV.

604

605 *Description*. Shell small to medium sized, suborbicular, slightly longer than high, strongly inflated,  
 606 equilateral; dorsal margin short and straight; beak acute, orthogyrate and slightly projecting above  
 607 dorsal margin; auricles incompletely preserved but clearly demarcated from disc; incremental lines  
 608 on anterior auricle indicate presence of deep byssal notch; ctenolium absent; 18 strong radial plicae  
 609 equal in strength, with regularly arranged tubercles at top of steep crests, starting close to beak, ribs  
 610 on distal part of disc slightly curved, posterior auricle showing commarginal ribs, interspaces  
 611 between ribs broad and deep.

612

613 *Remarks*. *Chlamys*-like bivalves were diverse in the Middle and Late Triassic, but their generic  
 614 assignment is often doubtful. One example is represented by Bittner's (1902) *Pecten volaris*,  
 615 subsequently assigned to *Chlamys* by Allasinaz (1972). As noted by Bittner (1902, p. 635), this species  
 616 is unusual among morphologically similar forms from the Triassic for having a strongly vaulted RV.  
 617 Bittner (1902) concluded that this species might have given rise to 'Vola' (= *Pecten*) -like taxa in the

1  
2  
3 618 Jurassic, which have generally a RV that is stronger vaulted in comparison to the LV. However,  
4  
5 619 '*Pecten*' *volaris* lacks a ctenolium and therefore cannot be attributed to *Pecten* or *Chlamys*.  
6  
7  
8 620 The specimens from Dobrogea agree very well with all main features of *Pecten volaris* as described  
9  
10 621 by Bittner (1902) and revised by Allasinaz (1972). However, neither Bittner (1902) nor Allasinaz  
11  
12 622 (1972) mentioned regularly arranged tubercles at the top of the rib crests, which are also not visible  
13  
14  
15 623 on the figures provided by these authors. We ascribe this lack of tubercles to an incomplete  
16  
17 624 preservation.  
18  
19  
20 625  
21  
22  
23 626 *Ecology*. Epibyssate suspension-feeders that rested pleurothetically on their RVs.  
24  
25  
26 627  
27  
28  
29 628 *Occurrence*. Cassian Formation (Italy, Longobardian–Julian; Zardini 1981), Gajine by Čevljanovići  
30  
31 629 (Bosnia and Herzegovina, Julian; Bittner 1902; Allasinaz 1972).  
32  
33  
34  
35 630  
36  
37  
38 631 Aviculopectinoidea n. gen. et n. sp.  
39  
40 632 Figures 4E–G  
41  
42  
43 633  
44  
45  
46 634 *Material*. Two RVs and one LV.  
47  
48  
49 635  
50  
51  
52 636 *Description*. Shell medium sized, infracrescent, both valves strongly inflated, suborbicular, higher  
53  
54 637 than long, subequilateral; umbo orthogyrate; beak acute, located in the middle of the straight dorsal  
55  
56 638 margin and slightly projecting above it; auricles clearly demarcated from disc, posterior auricles  
57  
58 639 obtuse; RV (Figs 4E and 4G): anterior auricle elongated, acute, with deep byssal notch lacking a  
59  
60

640 ctenolium; LV (Fig. 4F): anterior auricle triangular with small byssal sinus; faint antimarginal lines and  
641 growth lines on disc, where preserved; anterior auricle externally covered with slightly irregular  
642 commarginal ribs; internal structures unknown.

643

644 *Remarks.* The strongly inflated RV resembles '*Pecten*' *volaris* described above, but the new taxon  
645 lacks the pronounced radial plicae that characterize the latter species. A very unusual feature is the  
646 strong and approximately equal inflation of both valves, because pectinoid and aviculopectinoid  
647 bivalves usually have differentially vaulted valves, with at least one valve being flatter than in this  
648 taxon. Because the LV of '*Pecten*' *volaris* is unknown, it is unclear whether the two taxa differ in this  
649 respect.

650

651 *Ecology.* The deep byssal notch suggests that this species was an epibyssate suspension-feeder, but  
652 the equally inflated valves contradict a pleurothetic mode of life, which is common in pectinoids and  
653 aviculopectinoids. We therefore assume an orthothetic mode of life for these specimens.

654

655 Family ANTIJANIRIDAE Hautmann in Carter *et al.*, 2011

656 Genus AMPHIJANIRA Bittner, 1901

657

658 *Type species.* *Pecten janirula* Bittner, 1895.

659

660 *Amphijanira?* aff. *landrana* (Bittner, 1895)

661 Figures 4J–K

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

662  
  
663  
  
664  
  
665  
  
666  
  
667  
  
668  
  
669  
670  
671  
672  
673  
674  
675  
676  
677  
  
678  
  
679  
680  
681  
682  
  
683

aff. \*1895 *Pecten landranus*; Bittner, p. 166, pl. 19, fig. 21.

aff. 1972 *Amphijanira? landrana* (Bittner); Allasinaz, p. 269, pl. 33, figs 3–5 [cum syn.].

aff. 1981 *Amphijanira? landrana* (Bittner); Zardini, pl. 18, figs 4–5, 9a–b.

*Material.* Two incompletely preserved LVs and one questionable LV.

*Description.* Shell medium sized, infracrescent, equilateral, higher than long, weakly inflated; beak acute and orthogyrate; auricles not preserved; radial ribs with rounded crests, approximately 20 first order ribs starting close to beak, second order ribs irregularly intercalated, third and fourth order ribs occasionally present, similar in strength than second order ribs, higher order ribs closer to first order ribs than to second order ribs, interspaces between first order ribs broadest in the middle of disc where ribs of different orders can be easily distinguished from each other, at the anterior and posterior part of the disc interspaces between first order ribs smaller and second order ribs rarely intercalated, higher order ribs missing in these parts, fine and regular growth lines forming small nodes on the flanks of ribs where crossing them.

*Remarks.* The Dobrogea specimens might represent an ancestor of *Pecten landrana* Bittner, 1895, a Carnian bivalve revised by Allasinaz (1972). The sculpture of the LV of *P. landrana* is similar to the Dobrogea specimens. However, the ribs of the Dobrogea specimens are thicker. Allasinaz (1972) provisionally assigned *Pecten landrana* to the genus *Amphijanira*, which is followed herein.

684 *Ecology*. Species of the genus *Amphijanira* were probably epibyssate suspension-feeders (Ros-Franch  
685 *et al.* 2014).

686

687 Family HETEROPECTINIDAE Beurlen, 1954

688 Genus NEOMORPHOTIS Yin & Yin, 1983

689

690 *Type species. Neomorphotis gigantea* Yin & Yin, 1983.

691

692 *Neomorphotis comta* (Goldfuss, 1833 in Goldfuss 1833–1841)

693 Figures 4L–N

694

695 \*1833 *Ostrea comta*; Goldfuss, p. 4, pl. 72, fig. 6.

696 1835 *Spondylus comtus* (Goldfuss); Goldfuss, p. 93, pl. 105, fig. 1

697 1900 *Pseudomonotis beneckeii*; Bittner, p. 574, pl. 23, fig. 5.

698 1923 *Prospondylus comptus* (Goldfuss); Diener, p. 122 [cum syn.].

699 1926 *Prospondylus comptus* (Goldfuss); Reis, p. 124, pl. 9, figs 12–14.

700 1937 *Prospondylus comtus* (Goldfuss); Assmann, p. 58, pl. 12, fig. 1.

701 1967 *Prospondylus cf. comptus* (Goldfuss); Speciale, p. 1105, pl. 82, fig. 2 [cum syn.].

702 2008 *Neomorphotis compta* (Goldfuss); Posenato, p. 101, figs 4A–G [cum syn.].

703

1  
2  
3 704 *Material.* 22 LVs, one RV, two articulated specimens and probably six additional specimens with  
4  
5 705 unknown orientation. Specimens mostly incomplete.  
6  
7  
8 706  
9  
10  
11 707 *Description.* Shell large to very large, infracrescent, inequivalve, inequilateral, supposedly higher than  
12  
13 708 long, antero-dorsal margin shorter than postero-dorsal one; beaks prosogyrate; LV: inflated; beak  
14  
15 709 projecting above dorsal margin; anterior auricle smaller than posterior wing, obtuse, relatively well  
16  
17 710 demarcated from umbonal region, no byssal sinus; posterior wing faintly demarcated from umbonal  
18  
19 711 region, never completely preserved but probably obtuse; RV: less inflated than LV; beak not  
20  
21 712 projecting above dorsal margin; anterior auricle elongated, clearly demarcated from umbonal region  
22  
23 713 by deep byssal notch; LV possesses five to seven strong first order ribs with tubercles, starting close  
24  
25 714 to beak, second order ribs regularly intercalated in centre of first order interspaces, significantly  
26  
27 715 weaker than first order ribs, sometimes with small tubercles, four to six additional ribs of  
28  
29 716 subordinate order occur, ribs partly visible on internal moulds, fine growth lines occasionally  
30  
31 717 preserved, sculpture of RV generally stronger than in LV, first order ribs strongest in posterior region  
32  
33 718 and decreasing in strength anteriorly, hardly distinguishable in the middle and the anterior region  
34  
35 719 from higher order ribs, tubercles only visible on the strong first order ribs in the posterior region,  
36  
37 720 second order ribs starting closely to the beak, one to three third order ribs irregularly intercalated.  
38  
39  
40  
41  
42  
43 721  
44  
45  
46 722 *Remarks.* The original specific name *comta* introduced by Goldfuss (1833) was corrected by Frech  
47  
48 723 (1909) to *compta*, because Frech considered *comta* to be an inadvertent spelling error. However,  
49  
50 724 ‘comtus’ was commonly used in the sense of ‘ornate’ in Goldfuss’ time (e.g. Doederlein 1829, p. 259),  
51  
52 725 thus Goldfuss had chosen this word by intention and the correction of Frech (1909) is unjustified.  
53  
54 726 Posenato (2008) provided a detailed discussion of *Neomorphotis comta*, to which the reader is  
55  
56 727 referred.  
57  
58  
59  
60 728

729 *Ecology.* The byssal notch suggests that these bivalves were epibyssate suspension-feeders that  
730 rested pleurothetically on their RVs.

731

732 *Occurrence.* Jena Formation (Germany, late Bithynian – early Illyrian; Assmann 1937), Karlstadt  
733 Formation, Heilbronn Formation, Diemel Formation (Germany, early Illyrian; Assmann 1937), Dont  
734 Formation (Italy, Pelsonian–Illyrian; Posenato 2008), Valle Sabbia (Italy, middle Anisian; Speciale  
735 1967), Wetterstein Limestone (Austria, Ladinian; Reis 1926).

736

737 Family ASOELLIDAE Begg & Campbell, 1985

738 Genus ASOELLA Tokuyama, 1959

739

740 *Type species.* *Eumorphotis (Asoella) confertoradiata* Tokuyama, 1959.

741

742 *Asoella?* sp.

743 Figure 4I

744

745 *Material.* One incompletely preserved LV.

746

747 *Description.* Shell small, subcircular, slightly longer than high, moderately inflated; dorsal margin  
748 short, probably lacking well-differentiated auricles; beak small, obtuse, broadly rounded,  
749 orthogyrate, slightly projecting above dorsal margin; weak and regular growth lines, best visible at  
750 shell margins.



1  
2  
3 751  
4  
5  
6 752 *Remarks.* The subcircular shape, position of the beak, shell inflation, and the probable lack of  
7  
8 753 well-differentiated auricles in the LV are indicative for the genus *Asoella*. However, incomplete  
9  
10 754 preservation and the absence of information on the shell interior and the morphology of the RV  
11  
12  
13 755 make this assignment provisional.  
14  
15  
16 756  
17  
18  
19 757 *Ecology.* The shallow byssal notch in *Asoella* (Cox *et al.* 1969; not observed in the present material)  
20  
21 758 indicates an epibyssate mode of life.  
22  
23  
24 759  
25  
26  
27 760 Genus LEPTOCHONDRIA Bittner, 1891  
28  
29  
30 761  
31  
32  
33 762 *Type species.* *Pecten aeolicus* Bittner, 1891.  
34  
35  
36 763  
37  
38  
39 764 *Leptochondria pervulgata* (Bittner, 1902)  
40  
41 765 Figures 5A–B  
42  
43  
44 766  
45  
46  
47 767 \*1902 *Pecten* (?*Entolium*) *pervulgatus*; Bittner, p. 609, pl. 26, fig. 25.  
48  
49  
50 768 1972 *Leptochondria pervulgata* (Bittner); Allasinaz, p. 248, pl. 30, figs 2–4.  
51  
52  
53 769  
54  
55  
56 770 *Material.* 15 LVs and possibly two fragmented LVs.  
57  
58  
59 771  
60

772 *Description.* Shell medium sized, equilateral, subcircular, slightly longer than high, weakly inflated;  
 773 beak orthogyrate, positioned in the middle of straight dorsal margin and not projecting above it;  
 774 auricles faintly demarcated from disc, obtuse, sub-equal in size; approximately 20 to 25 widely  
 775 spaced ribs of first and second order with rounded crests, first order ribs starting close to the beak,  
 776 second order ribs slightly weaker, starting shortly below beak, regularly intercalated, few third order  
 777 ribs present, very weak and irregularly inserted by intercalation, ribs strongest in the middle part of  
 778 the disc and near the ventral margin, weaker at the anterior and posterior margins, auricles without  
 779 radial ribs, but fine and regular growth lines visible to the naked eye, most clearly on distal parts of  
 780 disc and auricles.

781

782 *Remarks.* The specimens from Dobrogea agree very well with all main features of *Leptochondria*  
 783 *pervulgata* as described by Bittner (1902) and revised by Allasinaz (1972). Bittner (1902) reported 25  
 784 to 30 ribs (or even more) that are irregularly distributed, resulting in an irregular pattern of stronger  
 785 and weaker ribs. This irregular pattern probably results from the irregularly intercalated third order  
 786 ribs. In contrast, Allasinaz (1972) described the thinner second order ribs to be regularly intercalated  
 787 between the first order ribs, but he did not mention any ribs of third order. However, the specimens  
 788 shown in Allasinaz (1972), which are the original specimens of Bittner (1902), show irregularly  
 789 intercalated third order ribs, which occur also in the Dobrogea specimens. Only one specimen shown  
 790 in Allasinaz (1972) possesses two completely preserved auricles that are sub-equal in size, which is  
 791 also the case in the Dobrogea specimens.

792

793 *Ecology.* Specimens of this genus have a flat RV with a byssal notch (not observed in the present  
 794 material), suggesting that these bivalves were epibyssate suspension-feeders that rested  
 795 pleurothetically on their RVs (Hautmann *et al.* 2013; Hofmann *et al.* 2014).

796

1		
2		
3	797	<i>Occurrence:</i> Clade near Čevljanovići (Bosnia and Herzegovina, Anisian; Bittner 1902), Monte Rite
4		
5	798	(Italy, Anisian; Allasinaz 1972).
6		
7		
8	799	
9		
10		
11	800	<i>Leptochondria separata</i> (Reis, 1926)
12		
13		
14	801	Figures 5C–D
15		
16		
17	802	
18		
19		
20	803	*1926 <i>Pecten separatus</i> ; Reis, p. 122, pl. 9, fig. 9.
21		
22		
23	804	1972 <i>Leptochondria separata</i> (Reis); Allasinaz, p. 251, pl. 30, fig. 8 [cum syn.].
24		
25		
26	805	
27		
28		
29	806	<i>Material.</i> Four LVs and one questionable LV.
30		
31		
32	807	
33		
34		
35	808	<i>Description.</i> Shell small, subcircular and slightly higher than long, weakly inflated, more or less
36		
37	809	equilateral; beak orthogyrate, positioned in the middle of straight dorsal margin and slightly
38		
39	810	projecting above it; auricles not well demarcated from disc, obtuse, probably equal in size, broad but
40		
41		
42	811	shallow byssal sinus below anterior auricle; up to 27 radial ribs on disc, intercalated in two ranks, first
43		
44	812	order ribs starting close to beak, increasing in strength towards the ventral margin, crests rounded,
45		
46	813	second order ribs irregularly intercalated, growth lines very weak, auricles occasionally ornamented
47		
48	814	with weak radial ribs and growth lines, growth lines most clearly developed on auricles.
49		
50		
51		
52	815	
53		
54		
55	816	<i>Remarks.</i> The lectotype, refigured in Allasinaz (1972), shows a broad but shallow byssal sinus, which
56		
57	817	is also present in the material from Dobrogea. <i>Leptochondria separata</i> differs from <i>Leptochondria</i>
58		
59		
60		

818 *pervulgata* by its smaller size, a more obtuse beak projecting above the dorsal margin, a higher  
 819 number of more densely spaced ribs and weaker growth lines.

820

821 *Ecology.* Same as for *Leptochondria pervulgata*.

822

823 *Occurrence.* Wetterstein Limestone (Austria, Ladinian; Reis 1926, Allasinaz 1972).

824

825 *Leptochondria viezzenensis* (Wilckens, 1909)

826 Figures 5F–H

827

828 \*1909 *Pecten viezzenensis*; Wilckens, p. 147. pl. 5, fig. 25.

829 1972 *Leptochondria viezzenensis* (Wilckens); Allasinaz, p. 259, pl. 31, figs 5–9.

830 2003 *Leptochondria cf. viezzenensis* (Wilckens); Szente & Vörös, p. 128, pl. Biv-II, figs 9–11.

831 2012 *Leptochondria viezzenensis* (Wilckens); Wasmer *et al.*, p. 1058, figs 7 (K–R).

832

833 *Material.* Four well preserved LVs and one poorly preserved LV.

834

835 *Description.* LV small to medium sized, infracrescent, subcircular, longer than high, equilateral,  
 836 weakly inflated; beak obtuse, orthogyrate, positioned in the middle of straight dorsal margin and  
 837 slightly projecting above it; auricles faintly demarcated from disc, anterior auricle obtuse and slightly  
 838 larger than posterior one, posterior auricle with shallow sinus; up to 40 ribs of three orders with

1  
2  
3 839 rounded crests, interspaces broader than ribs, thin first order ribs regularly arranged, starting close  
4  
5 840 to beak, second order ribs regularly intercalated, starting a few mm below the beak and almost as  
6  
7 841 broad as first order ribs, few third order ribs irregularly intercalated in the middle part of the disc,  
8  
9 842 ribs on auricles weaker developed than on the disc, fine and regularly growth lines visible on disc and  
10  
11 843 auricles.  
12  
13  
14  
15 844  
16  
17  
18 845 *Remarks.* The Dobrogea specimens match very well with the features of *Leptochondria viezzenensis*  
19  
20 846 described in Wilckens (1909), Allasinaz (1972) and Wasmer *et al.* (2012). However, Wilckens (1909)  
21  
22 847 mentioned that the growth lines on the auricles are more distinct than those on the disc, which is not  
23  
24 848 the case in the Dobrogea specimens. Furthermore, Allasinaz (1972) describes the ribs as irregularly  
25  
26 849 intercalated, which is only the case for the third order ribs of the Dobrogea specimen. However, in all  
27  
28 850 other points the characters of *L. viezzenensis* match those of the Dobrogea specimens. According to  
29  
30 851 Wasmer *et al.* (2012, p. 1059) also *Leptochondria minima* (Kiparisova, 1938) and *Leptochondria*  
31  
32 852 *virgalensis* (von Wittenburg, 1909) may possess a shallow sinus in the left posterior auricle, but the  
33  
34 853 sculptures of these species differ clearly from the sculpture of *L. viezzenensis*. In contrast to  
35  
36 854 *Leptochondria pervulgata*, which is also longer than high, *L. viezzenensis* possesses a beak that is  
37  
38 855 slightly projecting above the dorsal margin, ribbed auricles, a posterior auricle with a shallow sinus  
39  
40 856 and a higher number of ribs.  
41  
42  
43  
44  
45  
46 857  
47  
48  
49 858 *Ecology.* Same as for all other *Leptochondria* species.  
50  
51  
52 859  
53  
54  
55 860 *Occurrence.* Mianwali Formation (Pakistan, late Smithian – middle Spathian; Wasmer *et al.* 2012),  
56  
57 861 Felsőőrs Formation (Hungary, Pelsonian–Illyrian; Sente & Vörös 2003), Cima Viezzena (Italy,  
58  
59 862 Longobardian; Wilckens 1909; Allasinaz 1972).  
60

863

864

Family ORNITHOPECTINIDAE Hautmann in Carter *et al.*, 2011

865

Genus ORNITHOPECTEN Cox, 1962

866

867 *Type species. Aviculopecten bosniae* Bittner, 1902.

868

869

*Ornithopecten interruptus* (Bittner, 1902)

870

Figure 5l

871

872 \*1902 *Aviculopecten interruptus*; Bittner, p. 591, pl. 26, fig. 13.873 1972 *Ornithopecten interruptus* (Bittner); Allasinaz, p. 263, pl. 32, fig. 5.

874

875 *Material.* One well preserved LV.

876

877 *Description.* Shell medium sized, infracrescent, longer than high, inequilateral, weakly inflated; beak  
878 acute, orthogyrate, slightly projecting above straight dorsal margin, postero-dorsal margin more than  
879 twice as long as antero-dorsal margin; posterior wing elongated and flat, anterior auricle smaller  
880 than posterior wing, growth lines on wing and auricle suggestive of acute distal ends, shallow byssal  
881 sinus below anterior wing; approximately 22 rounded ribs that are straight on the posterior part of  
882 the shell but curved on the anterior part, approximately 10 first order ribs starting close to beak,  
883 separated by broad interspaces, one first order rib coinciding with the postero-dorsal margin, second

1  
2  
3 884 order ribs starting at various distances from the beak, irregularly intercalated, regular growth lines  
4  
5 885 reflected by slight elevations on shell.  
6  
7  
8 886  
9  
10  
11 887 *Remarks.* Bittner (1902) and Allasinaz (1972) describe *Ornithopecten interruptus* as medium sized,  
12  
13 888 suborbicular and weakly inflated bivalve that is longer than high and has an elongated posterior wing  
14  
15 889 that is larger than the anterior auricle. Both structures have acute distal ends and are poorly  
16  
17 890 demarcated from the disc. The Dobrogea specimen is in accordance with all these characters and  
18  
19 891 also shows high similarities to the sculpture of *O. interruptus*. Bittner (1902) reported six growth  
20  
21 892 disruptions that interrupt and shift the radial ribs. We regard this feature as a response to  
22  
23 893 environmental changes rather than a character of taxonomic importance. We follow Cox (1962) and  
24  
25 894 assign this species to the genus *Ornithopecten*.  
26  
27  
28  
29  
30 895  
31  
32  
33 896 *Ecology.* Specimens of the genus *Ornithopecten* have a less inflated RV with a byssal notch (not  
34  
35 897 observed in the present material), suggesting that these bivalves were epibyssate suspension-feeders  
36  
37 898 that rested pleurothetically on their RVs.  
38  
39  
40  
41 899  
42  
43  
44 900 *Occurrence.* Clade near Čevljanovići (Bosnia and Herzegovina, Anisian; Bittner 1902; Allasinaz 1972).  
45  
46  
47 901  
48  
49  
50 902 *Ornithopecten cf. katzeri* (Bittner, 1902)  
51  
52 903 Figure 5E  
53  
54  
55 904  
56  
57  
58 905 cf. \*1902 *Aviculopecten katzeri*; Bittner, p. 608, pl. 26, fig. 22.  
59  
60

906 cf. 1972 *Ornithopecten katzeri* (Bittner); Allasinaz, p. 263, pl. 32, fig. 4.

907

908 *Material.* One almost completely LV.

909

910 *Description.* Shell small, ovate, inequilateral, weakly inflated; beak prosogyrate, acute; antero-dorsal  
911 margin straight and probably longer than postero-dorsal margin; wings faintly demarcated from disc,  
912 anterior wing small; seven first order ribs, starting close to beak, generally separated by broad  
913 interspaces, ribs slightly curved, rib interspaces on anterior part narrower than on posterior part, up  
914 to five thin second order ribs intercalated between first order ribs, weak commarginal ribs forming  
915 tubercles where they cross first order ribs.

916

917 *Remarks.* The Dobrogea specimen is very similar to *Ornithopecten katzeri*, which, however, is based  
918 on a single RV, which makes the assignment of the present LV uncertain. Bittner (1902) and Allasinaz  
919 (1972) mentioned a slightly higher number of ribs (ten first order ribs and up to eight second order  
920 ribs between first order ribs), which might be due to the slightly larger size of the type specimen. *O.*  
921 cf. *katzeri* can be distinguished from *Ornithopecten interruptus* by the prosogyrate beak, the lower  
922 number of first order ribs and the higher number of second order ribs.

923

924 *Ecology.* Same as for *Ornithopecten interruptus*.

925

926 *Occurrence.* Clade near Čevljanovići (Bosnia and Herzegovina, Anisian; Bittner 1902; Allasinaz 1972).

927



1		
2		
3	928	Superfamily PECTINOIDEA Rafinesque, 1815
4		
5		
6	929	Family PECTINIDAE Rafinesque, 1815
7		
8		
9	930	Genus AVICHLAMYS Allasinaz, 1972
10		
11	931	
12		
13		
14	932	<i>Type species. Pecten csopakensis</i> Frech, 1905.
15		
16		
17	933	
18		
19		
20	934	<i>Avichlamys?</i> cf. <i>nicolensis</i> (Ogilvie Gordon, 1927)
21		
22		
23	935	Figure 50
24		
25		
26	936	
27		
28		
29	937	cf. *1927 <i>Pecten nicolensis</i> ; Ogilvie Gordon, p. 26, pl. 2, fig. 22.
30		
31		
32	938	cf. 1972 <i>Avichlamys nicolensis</i> (Ogilvie Gordon); Allasinaz, p. 370, pl. 48, fig. 11.
33		
34		
35	939	
36		
37		
38	940	<i>Material.</i> One incompletely preserved LV.
39		
40		
41	941	
42		
43		
44	942	<i>Description.</i> Shell medium sized, infracrescent, almost flat; postero-dorsal part not preserved;
45		
46	943	anterior auricle incompletely preserved, well demarcated from disc; at least 16 distinct, narrow but
47		
48	944	high radial first order ribs with rounded crests, wide interspaces getting broader distally where thin
49		
50	945	second order ribs are intercalated, second order ribs positioned asymmetrically in-between first
51		
52	946	order ribs, commarginal riblets also present, creating a reticular pattern by crossing radial ribs,
53		
54	947	distance between commarginal riblets decreasing towards ventral margin.
55		
56		
57		
58	948	
59		
60		

949 *Remarks.* Two Triassic pectinoid species have an ornamentation similar to the Dobrogea specimen,  
 950 namely *Pecten reticulatus* von Schlotheim, 1820 and *Avichlamys nicolensis* (Ogilvie Gordon, 1927),  
 951 but the latter is weaker inflated (Ogilvie Gordon 1927). Because the Dobrogea specimen is almost  
 952 flat, we provisionally assign it to *A. nicolensis*. If correct, this implies that *Avichlamys* survived into the  
 953 Middle Triassic.

954

955 *Ecology.* Epibyssate suspension-feeder that rested pleurothetically on its RV.

956

957 *Occurrence.* Werfen Formation (Italy, Early Triassic; Ogilvie Gordon 1927; Allasinaz 1972).

958

959 Genus CHLAMYS Röding in Bolten, 1798

960

961 *Type species.* *Pecten islandicus* Müller, 1776.

962

963 Subgenus PRAECHLAMYS Allasinaz, 1972

964

965 *Type species.* *Pecten (Chlamys) inaequalternans* Parona, 1889.

966

967 *Chlamys (Praechlamys) prima* n. sp.

968 Figures 5J–M

969

970 *LSID.* urn:lsid:zoobank.org:act:6EDFD317-06D9-4492-BF45-ED9A60654390

1  
2  
3 971  
4  
5  
6 972 *Derivation of name. Prima* (Latin: the first) because this species is among the geologically earliest  
7  
8 973 species of its genus.  
9  
10  
11 974  
12  
13  
14 975 *Holotype*. LPB IIL 2062  
15  
16  
17 976 *Paratypes*. LPB IIL 2060, 2061, 2063, 2064, 2065  
18  
19  
20  
21 977  
22  
23  
24 978 *Material*. Twelve fragmented but otherwise well preserved LVs, two internal moulds of LVs with  
25  
26 979 some shell remains, one external mould of a LV and one unassigned valve.  
27  
28  
29 980  
30  
31  
32 981 *Diagnosis. Praechlamys* with approximately 30 radial ribs arranged in two ranks on the LV; anterior  
33  
34 982 disc margin demarcated from anterior auricle by steep slope; shallow radial depression on posterior  
35  
36 983 part of disc, where radial ornament becomes weaker than on main part of disc.  
37  
38  
39  
40 984  
41  
42  
43 985 *Description*. Shell medium sized, moderately inflated, inequilateral, suborbicular, approximately as  
44  
45 986 high as long, disc inflation decreases towards the posterior margin, separating morphologically the  
46  
47 987 posterior part from the remaining disc; beak slightly prosogyrate, positioned in the middle of the  
48  
49 988 straight dorsal margin and slightly projecting above it; auricles relatively large, anterior auricle with  
50  
51 989 byssal sinus and larger than posterior one, clearly demarcated from disc by steep slope, posterior  
52  
53 990 auricle truncated, obtuse and smaller than anterior one, clearly demarcated from disc but slope not  
54  
55 991 as steep as anteriorly; up to 30 rounded first and second order ribs bearing small tubercles, getting  
56  
57 992 broader towards the ventral margin, first order ribs starting close to beak, second order ribs starting  
58  
59 993 slightly more distally, regularly intercalated, ribs denser but weaker on flat posterior part of disc,  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

994 growth lines present, two weak ribs and several riblets on posterior auricle; internal structures not  
995 observed.

996  
997 *Remarks.* The most similar species is *Chlamys (Praechlamys) subdivisa* (Bittner, 1901), which differs in  
998 having a tripartite division of the disc with a broad middle part and narrow radial depressions  
999 anteriorly and posteriorly. In the Dobrogea specimens a comparable depression/flattening of the disc  
1000 occurs only posteriorly. Furthermore, *C. (P.) subdivisa* has 40 to 50 smooth ribs while *C. (P.) prima*  
1001 has only approximately 30 ribs. The intercalation of second order ribs in *C. (P.) subdivisa* is more  
1002 irregular and the posterior auricle lacks ribs or riblets.

1003  
1004 *Ecology.* Specimens of *Chlamys (Praechlamys)* have a slightly less inflated RV with a deep byssal  
1005 notch below the anterior auricle (as inferred from the presence of a byssal sinus in the LV). This  
1006 suggests that these bivalves were epibyssate suspension-feeders that rested pleurothetically on their  
1007 RVs.

1008

1009 Genus UNCERTAIN

1010

1011 Pectinidae indet.

1012 Figures 5P–Q

1013

1014 *Material.* One LV and one RV, both incompletely preserved internal moulds.

1015

1  
2  
3 1016 *Description.* Shell small, suborbicular, subequilateral, slightly higher than long; LV weakly inflated, RV  
4  
5 1017 less inflated than LV; approximately ten regularly arranged, broad radial folds with narrow  
6  
7 1018 interspaces, irregular growth lines.  
8  
9  
10  
11 1019  
12  
13  
14 1020 *Remarks.* These specimens are unlike all other pectinoids from Dobrogea with respect to their  
15  
16 1021 ornamentation. Because the present specimens are internal moulds, we cannot reconstruct the  
17  
18 1022 original ornament in detail, except from the fact that the species had ten main plicae. We therefore  
19  
20 1023 present this species in open nomenclature.  
21  
22  
23  
24 1024  
25  
26  
27 1025 *Ecology.* The less inflated RV suggests that these bivalves were epibyssate suspension-feeders that  
28  
29 1026 rested pleurothetically on their RVs.  
30  
31  
32 1027  
33  
34  
35 1028 Family PLEURONECTITIDAE Hautmann in Carter *et al.*, 2011  
36  
37  
38 1029 Genus PLEURONECTITES von Schlotheim, 1820  
39  
40 1030  
41  
42  
43 1031 *Type species.* *Pleuronectites laevigatus* von Schlotheim, 1820.  
44  
45  
46 1032  
47  
48  
49 1033 *Pleuronectites?* cf. *balatonicus* (Bittner, 1901)  
50  
51  
52 1034 Figure 5N  
53  
54  
55 1035  
56  
57  
58 1036 cf. \*1901 *Pecten balatonicus*; Bittner, p. 37, pl. 5, figs 9–11.  
59  
60 1037 ? 1972 *Pleuronectites balatonicus* (Bittner); Allasinaz, p. 323, pl. 43, fig. 5 (non fig. 6).

1038

1039 *Material.* One almost complete RV and possibly one additional, incompletely preserved RV.

1040

1041 *Description.* Shell of RV small to medium sized, weakly inflated or nearly flat, inequilateral, higher  
 1042 than long; beak orthogyrate, positioned in the middle of straight dorsal margin and not projecting  
 1043 above it; auricles unequal in size, anterior auricle elongated, clearly demarcated from disc by deep  
 1044 byssal notch, posterior auricle obtuse, triangular, clearly demarcated from disc; shell exterior smooth  
 1045 except for weak, irregularly spaced incremental lines and weak radial striae that possibly represent  
 1046 radial sectors of the shell microstructure, as described by Carter & Hautmann (2011).

1047

1048 *Remarks.* The shape of this specimen is virtually identical to that of Bittner's (1901) type specimen of  
 1049 *Pecten balatonicus*, except for a relatively shorter anterior auricle. However, it seems possible that  
 1050 this difference is due to incomplete preservation. Although we provisionally follow Allasinaz' (1972)  
 1051 assignment to *Pleuronectites*, we note that this assignment is problematic. First, the presence of a  
 1052 ctenolium cannot be demonstrated, although this feature might be absent due to incomplete  
 1053 preservation. Second, the ligament morphology is unknown. The specimens from the Cassian  
 1054 Formation that Allasinaz (1972, pl. 34, fig. 6a–c) assigned to *P. balatonicus* do clearly not belong to  
 1055 *Pleuronectites* (see revision of *Pleuronectites* in Hautmann 2010) but to an entoliid species. Still, the  
 1056 overall morphology, smooth shell exterior and the radial arrangement of shell-microstructural  
 1057 features are suggestive for *Pleuronectites* (Hautmann 2010; Carter & Hautmann 2011). The outline of  
 1058 this specimen is also very similar to the new aviculoid taxon described above (Figs 4E–G), but its valve  
 1059 is nearly flat, which is not due to compaction because no deformation features are observed.

1060

1		
2		
3	1061	<i>Ecology.</i> The byssal notch in the RV suggests that this species was an epibyssate suspension-feeder
4		
5	1062	that rested pleurothetically on its RVs (Hautmann 2010).
6		
7		
8	1063	
9		
10		
11	1064	<i>Occurrence.</i> Veszprémer Mergel (Hungary, Carnian; Bittner 1901; Allasinaz 1972).
12		
13		
14	1065	
15		
16		
17	1066	Suborder ENTOLIIDINA Hautmann in Carter <i>et al.</i> , 2011
18		
19		
20	1067	Superfamily ENTOLIOIDEA von Teppner, 1922
21		
22		
23		
24	1068	Family ENTOLIIDAE von Teppner, 1922
25		
26		
27	1069	Genus ENTOLIUM Meek, 1865
28		
29	1070	
30		
31		
32	1071	<i>Type species.</i> <i>Pecten demissus</i> Meek, 1865.
33		
34		
35	1072	
36		
37		
38	1073	<i>Entolium reticulatum</i> n. sp.
39		
40	1074	Figure 5R
41		
42		
43	1075	
44		
45		
46	1076	<i>LSID.</i> urn:lsid:zoobank.org:act:CADEA6B4-C9F9-41C0-8D51-BF6F62E5BA95
47		
48		
49	1077	
50		
51		
52	1078	<i>Derivation of name.</i> <i>Reticulum</i> (Latin: net), referring to the reticular sculpture.
53		
54		
55	1079	
56		
57		
58	1080	<i>Holotype.</i> LPB IIL 2070
59		
60		

- 1081
- 1082 *Material.* Two LVs.
- 1083
- 1084 *Diagnosis.* *Entolium* with weak reticular ornamentation, resulting from thin and regular radial folds
- 1085 crossing commarginal folds.
- 1086
- 1087 *Description.* Shell small, disc weakly inflated, subcircular, equilateral; beak orthogyrate, located in the
- 1088 middle of straight dorsal margin and projecting above it, two weak depressions located at the
- 1089 anterior and posterior part of the disc, probably corresponding to internal lateral disc buttresses;
- 1090 auricles obtuse and equal in size, posterior auricle demarcated more clearly from disc than anterior
- 1091 one, growth lines present, no byssal sinus; thin and regular radial folds, most clearly developed on
- 1092 umbonal region, regular commarginal folds, almost as broad or slightly broader than radial folds,
- 1093 most clearly on umbonal region, radial and commarginal folds form weak reticular pattern; internal
- 1094 characters unknown.
- 1095
- 1096 *Remarks.* The ornamentation of this specimen is different from all Triassic entoliids known to us.
- 1097
- 1098 *Ecology.* Specimens of the genus *Entolium* lack a byssal notch, suggesting that these bivalves where
- 1099 unattached epifaunal suspension feeders.
- 1100
- 1101 *Entolium* sp.
- 1102 Figure 5V
- 1103



1		
2		
3	1104	<i>Material.</i> Two LVs.
4		
5		
6	1105	
7		
8		
9	1106	<i>Description.</i> Shell small, weakly inflated, inequilateral, suborbicular, higher than long; beak
10		
11	1107	orthogyrate and obtuse; two radial depressions on dorsal part of disc probably corresponding to
12		
13	1108	internal disc buttresses, posterior despression slightly more developed than anterior one; auricles
14		
15	1109	clearly demarcated from disc, obtuse; growth lines and weak radial lines, visible on disc and auricles
16		
17		
18		
19	1110	
20		
21		
22	1111	<i>Remarks.</i> The Dobrogea specimens show similarities with <i>Entolium discites</i> (von Schlotheim, 1820)
23		
24	1112	and <i>Entolium hallense</i> (von Wöhrmann, 1889) as described in Allasinaz (1972). However, the
25		
26	1113	preservation state of the specimens, especially the poorly preserved auricles, prevent a clear
27		
28	1114	assignment to one of these two species. Therefore we present the Dobrogea specimens in open
29		
30	1115	nomenclature. <i>Entolium</i> sp. differs from <i>Entolium reticulatum</i> by the lack of the reticular
31		
32	1116	ornamentation.
33		
34		
35		
36	1117	
37		
38		
39	1118	<i>Ecology.</i> Same as for <i>Entolium reticulatum</i> .
40		
41		
42	1119	
43		
44		
45	1120	
46		
47		
48	1121	
49		
50		
51	1122	Genus ENTOLIOIDES Allasinaz, 1972
52		
53	1123	
54		
55		
56		
57	1124	<i>Type species.</i> <i>Pecten zitteli</i> von Wöhrmann & Koken, 1892.
58		
59		
60	1125	

1126 *Entolioides deecke* (Parona, 1889)

1127 Figures 5S–U

1128

1129 \*1889 *Pecten* (*Vola*?) *deecke*; Parona, p. 92, pl. 6, fig. 8.

1130 1892 *Pecten deecke* Parona; von Wöhrmann & Koken, p. 172, pl. 9, figs 1–2.

1131 1923 *Pecten* (*Entolium*) *deecke* Parona; Diener, p. 69 [cum syn.].

1132 1972 *Entolioides deecke* (Parona); Allasinaz, p. 297, pl. 37, figs 6–9 [cum syn.].

1133

1134 *Material*. Four LVs that are mostly incomplete and/or internal moulds with some shell remains.

1135 Possibly one additional LV and one incompletely preserved LV showing some details of the internal  
1136 morphology.

1137

1138 *Description*. Shell large, disc longer than high, subequilateral, weakly inflated; beak obtuse,  
1139 orthogyrate, not projecting above straight dorsal margin; depression subparallel to postero-dorsal  
1140 margin starting close to beak occasionally present (Fig. 5S), probably reflecting internal structural  
1141 elements; auricles obtuse, posterior auricle better demarcated from disc than anterior one, no byssal  
1142 sinus; first and second order ribs with slightly variable interspaces, approximately 11 first order ribs  
1143 starting close to beak and getting broader towards posterior margin, regularly intercalated second  
1144 order ribs starting somewhat below the beak, fine, dense and regularly arranged growth lines  
1145 producing irregular nodes and wavy crests on ribs, auricles with growth lines only; monomyarian,  
1146 adductor muscle scar rounded and located on the postero-ventral part of the shell (Fig. 5U), ligament  
1147 alivincular-alate, auricular crura present along lower border of auricles starting close to the resilifer,  
1148 growth lines and first order ribs present.

1		
2		
3	1149	
4		
5		
6	1150	<i>Remarks.</i> We assign the Dobrogea specimens to <i>Entolioides deecke</i> i based on size, shape and the
7		
8	1151	diagnostic ornamentation.
9		
10		
11	1152	
12		
13		
14	1153	<i>Ecology.</i> The byssal sinus of <i>Entolioides</i> (not preserved in the present material) is either weak or
15		
16	1154	lacking and the RV is less inflated than the LV (Allasinaz 1972, table 3). This suggests that species of
17		
18	1155	this genus were unattached epifaunal suspension feeders that rested pleurothetically on their RVs.
19		
20		
21		
22	1156	
23		
24		
25	1157	<i>Occurrence.</i> Schlernplateau beds (Italy, late Longobardian – early Julian; von Wöhrmann & Koken
26		
27	1158	1892; Allasinaz 1972), Acquate (Italy, Julian; Parona 1889; Allasinaz 1972), Valletta di Rogno (Italy,
28		
29	1159	Carnian; Allasinaz 1972).
30		
31		
32		
33	1160	
34		
35		
36	1161	Genus SCYTHENTOLIUM Allasinaz, 1972
37		
38	1162	
39		
40		
41	1163	<i>Type species.</i> <i>Pecten tirolicus</i> von Wittenburg, 1908.
42		
43		
44	1164	
45		
46		
47	1165	<i>Scythentolium anisicum</i> n. sp.
48		
49	1166	Figures 6A–D
50		
51		
52	1167	
53		
54		
55	1168	<i>LSID.</i> urn:lsid:zoobank.org:act:FA7D2579-A447-47EE-A277-732EB85105A0
56		
57		
58		
59	1169	
60		

1170 *Derivation of name.* Referring to the stratigraphic age of this species.

1171

1172 *Holotype.* LPB IIL 2075

1173 *Paratypes.* LPB IIL 2076, 2077, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2158

1174

1175 *Material.* Eight LVs, two RVs, three articulated and four fragmentary shells.

1176

1177 *Diagnosis.* *Scythentolium* with slightly prorescent disc.

1178

1179 *Description.* Shell large, disc subcircular, inequivalve, slightly prorescent; inflation of valves  
 1180 moderate, comparable in LV and RV; beak obtuse, orthogyrate and positioned in the middle of dorsal  
 1181 margin but not projecting above it; dorsal margin straight in LV and angulate in RV; auricles of LV  
 1182 subequal in size, obtuse, clearly demarcated from disc; auricles of RV obtuse, triangular, clearly  
 1183 demarcated from disc, byssal notch below anterior auricle; both valves smooth except for regular  
 1184 and fine growth lines and very fine antimarginal structure (Fig. 6D), growth lines and antimarginal  
 1185 lines also observable on auricles.

1186

1187 *Remarks.* A deep byssal notch and a smooth shell surface are characteristic for the genus  
 1188 *Scythentolium* Allasinaz, 1972. Because the Dobrogea specimens differ from all other Triassic  
 1189 *Scythentolium* species by the slightly prorescent growth, we erect a new species, *S. anisicum*. Our  
 1190 finds indicate that *Scythentolium* was not restricted to the Early Triassic but persisted into the Middle  
 1191 Triassic.

1		
2		
3	1192	
4		
5		
6	1193	<i>Ecology.</i> The byssal notch of <i>Scythentolium</i> suggests that species of this genus were epibyssate
7		
8	1194	suspension-feeders that rested pleurothetically on their RVs.
9		
10		
11	1195	
12		
13		
14	1196	Suborder OSTREINA Férussac, 1822 in Férussac 1821–1822
15		
16		
17	1197	Superfamily OSTREOIDEA Rafinesque, 1815
18		
19		
20	1198	Family OSTREIDAE Rafinesque, 1815
21		
22		
23	1199	Genus UMBROSTREA Hautmann, 2001 <i>b</i>
24		
25		
26	1200	
27		
28		
29	1201	<i>Type species.</i> <i>Umbrostrea emamii</i> Hautmann, 2001 <i>b</i> .
30		
31		
32	1202	
33		
34		
35	1203	<i>Umbrostrea?</i> sp.
36		
37	1204	Figure 6K
38		
39		
40	1205	
41		
42		
43	1206	<i>Material.</i> One internal mould of a LV.
44		
45		
46	1207	
47		
48		
49	1208	<i>Description.</i> Shell small, subelliptical, retrocrescent, with relatively large attachment surface and
50		
51	1209	strongly plicate shell margin; plicae rounded, regularly arranged but varying in thickness.
52		
53		
54	1210	
55		
56		
57	1211	<i>Remarks.</i> The curvature suggests that this specimen is a LV, which, in combination with the presence
58		
59	1212	of an attachment surface, leads to its identification as a ‘true’ oyster.
60		

1213

1214 *Ecology.* Cemented epifaunal suspension-feeder.

1215

1216 Suborder MONOTIDINA Waterhouse, 2001 (*sensu* Wasmer *et al.* 2012)

1217 Superfamily MONOTOIDEA Fischer, 1887 in Fischer 1880–1887

1218 Family MONOTIDAE Fischer, 1887 in Fischer 1880–1887

1219 Genus UNCERTAIN

1220

1221 Monotidae indet.

1222 Figures 6O–P

1223

1224 *Material.* One incompletely preserved internal mould of a LV.

1225

1226 *Description.* Shell medium sized, elliptical, longer than high, moderately inflated, inequilateral,

1227 retrocrescent; beak acute, slightly prosogyrate, slightly projecting above straight and relatively short

1228 dorsal margin; anterior and posterior wing weakly demarcated from disc; irregular and uneven radial

1229 ribs, increasing in number by intercalation, first order ribs starting close to beak, at the posterior part

1230 of the shell hardly distinguishable from second order ribs, second order ribs starting approximately in

1231 the middle part of the shell, generally narrow interspaces between ribs, interspaces on postero-

1232 dorsal part of shell are narrowest.

1233

1234 *Remarks.* *Monotis* is an index fossil for the middle Norian to early Rhaetian (McRoberts *et al.* 2008)  
1235 and likely arose from *Meleagrinnella* Whitfield, 1885 during the early or middle Norian (McRoberts  
1236 2010, 2017). It is therefore surprising to find a bivalve with a *Monotis*-like morphology in Anisian  
1237 strata. Given the unusual stratigraphic occurrence and the incomplete preservation, it seems possible  
1238 that this species represents a Middle Triassic homeomorph of *Monotis*. We therefore assign this  
1239 taxon only tentatively to the Monotidae.

1240

1241     *Ecology.* For *Monotis*, the type genus of the Monotidae, several modes of life have been suggested,

1242     ranging from epibyssally attached to hard substratum (Stanley 1972; Hallam 1981; McRoberts *et al.*

1243     2008) or plants (Grant-Mackie 1980; Ando 1987) to pseudoplanktonic (Hayami 1969; Stanley 1972;

1244     Silberling *et al.* 1997) or nektonic (Jefferies & Minton 1965). The exact mode of life of the Dobrogea

1245     specimen remains speculative.

1247 Suborder LIMINA Waller, 1978

1248 Superfamily LIMOIDEA Rafinesque, 1815

1249 Family LIMIDAE Rafinesque, 1815

1250 Genus MYSIDIOPTERA Salomon, 1895

1252     *Type species. Mysidioptera ornata* Salomon, 1895.

1254 *Mysidioptera fornicata* Bittner, 1895

1255 Figures 6E–G

1256

1257 \*1895 *Mysidioptera fornicata*; Bittner, p. 184, pl. 20, fig. 11.1258 2004 *Mysidioptera fornicata* Bittner; Komatsu *et al.*, fig. 7(2).

1259

1260 *Material.* 13 LVs, 15 RVs and two unassigned valves, one of them showing the shell's interior.

1261

1262 *Description.* Shell medium sized, moderately to strongly inflated, equivalve, inequilateral, slightly  
 1263 procrenscent, suborbicular to obliquely ovate in outline, height/length-ratio variable, either higher  
 1264 than long or vice versa; beak obtuse, prosogyrate; anterior margin concave, lunula deeply impressed,  
 1265 with byssal gape between valves, postero-dorsal margin straight; anterior auricle absent, posterior  
 1266 auricle obtuse, not well demarcated from main body of shell; externally smooth except for slightly  
 1267 irregular growth lines on disc and posterior auricle, occasionally fine radial ribs present on disc,  
 1268 hardly recognizable on umbonal region, lunula with strong growth lines.

1269

1270 *Remarks.* Nearly 100 *Mysidioptera* species have been reported from the Triassic and several of these  
 1271 are smooth or weakly ornamented (Posenato 2008). *Mysidioptera* species that are weakly  
 1272 ornamented with radial elements and strongly inflated include *M. fornicata* Bittner, 1895, *M. globosa*  
 1273 Broili, 1903, *M. inflata* Wilckens, 1909, *M. kittlii* Bittner, 1895 and *M. tommasii* Sacchi Vialli & Vai,  
 1274 1958. The Dobrogea specimens show most similarities with *M. fornicata*, which is characterized by a  
 1275 suborbicular outline, a higher than long shell, a straight postero-dorsal margin and the presence of  
 1276 radial striae on the lunula and in the umbonal region. According to Broili (1903), *M. globosa* has  
 1277 densely spaced, weak radial ribs that undulate where intersecting strong growth lines, a feature that  
 1278 is not present in the Dobrogea specimens. *M. kittlii* and *M. inflata* have the same ornamentation as



1  
2  
3 1279 *M. globosa* and Wilckens (1909) assigned these three species to the subgenus *Latemaria*, because of  
4  
5 1280 their strong inflation. Furthermore, *M. kittlii* does not possess a lunula or a byssal gape (Bittner  
6  
7 1281 1895). The beak of *M. tommasii* described by Speciale (1967) is acute and the inflation of the disc  
8  
9 1282 decreases towards the shell margin, which is not the case in the Dobrogea specimens.  
10  
11  
12  
13 1283  
14  
15  
16 1284 *Ecology*. The equivalved condition and the presence of a byssal notch suggest that specimens of *M.*  
17  
18 1285 *fornicata* rested orthothetically and byssally attached on the sediment.  
19  
20  
21 1286  
22  
23  
24 1287 *Occurrence*. Leidapo Member (China, late Pelsonian – early Illyrian; Komatsu *et al.* 2004), Cassian  
25  
26 1288 Formation (Italy, Longobardian–Julian; Bittner 1895).  
27  
28  
29  
30 1289  
31  
32  
33 1290 *Mysidioptera* sp.  
34  
35 1291 Figures 6L  
36  
37  
38 1292  
39  
40  
41 1293 *Material*. One articulated specimen and possibly one additional LV.  
42  
43  
44 1294  
45  
46  
47 1295 *Description*. Shell medium sized, moderately inflated, procrescent, equivalve, inequilateral,  
48  
49 1296 subtrigonal, higher than long; beak orthogyrate, acute, slightly projecting above probably straight  
50  
51 1297 dorsal margin; anterior margin concave; anterior auricles not preserved, posterior auricle small,  
52  
53 1298 obtuse, well demarcated from body shell; shell smooth except for fine and regular growth lines on  
54  
55 1299 body shell.  
56  
57  
58  
59 1300  
60

1301 *Remarks.* The overall morphology indicates that these specimens belong to the genus *Mysidioptera*.  
 1302 However, none of the smooth *Mysidioptera* species reported from the Triassic possesses a posterior  
 1303 auricle that is as well demarcated from the body shell as that of these specimens. Because of the  
 1304 limited material and incomplete preservation, we describe these specimens in open nomenclature  
 1305 rather than establishing a new species.

1306

1307 *Ecology.* Same as for *Mysidioptera fornicata*.

1308

1309 Genus PLAGIOSTOMA J. Sowerby, 1814

1310

1311 *Type species.* *Plagiostoma gigantea* J. Sowerby, 1814.

1312

1313 *Plagiostoma striatum* (von Schlotheim, 1820)

1314 Figures 6H–J

1315

1316 \*1820 *Chamites striatus*; von Schlotheim, p. 210.

1317 1823 *Chamites striatus* von Schlotheim; Schlotheim, pl. 34, fig. 1.

1318 1923 *Lima (Plagiostoma) striata* (von Schlotheim); Diener, p. 107 [cum syn.].

1319 1928 *Lima striata* (von Schlotheim); Schmidt, p. 154, fig. 326.

1320 1937 *Lima striata* (von Schlotheim); Assmann, p. 49, pl. 10, fig. 14.

1321 1967 *Plagiostoma striatum* (von Schlotheim); Speciale, p. 1109, pl. 82, fig. 7 [cum syn.].

1		
2		
3	1322	2003 <i>Plagiostoma striatum</i> (von Schlotheim); Sente & Vörös, p. 128, pl. 2, fig. 2 [cum syn.].
4		
5		
6	1323	2004 <i>Plagiostoma striatum</i> (von Schlotheim); Komatsu <i>et al.</i> , fig. 7(18).
7		
8		
9	1324	2008 <i>Plagiostoma striatum</i> (von Schlotheim); Posenato, p. 101, fig. 3O [cum syn.].
10		
11		
12	1325	2015 <i>Plagiostoma striatum</i> (von Schlotheim); Hagdorn, fig. 7.11.
13		
14		
15	1326	2016 <i>Plagiostoma striatum</i> (von Schlotheim); Brom <i>et al.</i> , fig 1E.
16		
17		
18	1327	2017 <i>Plagiostoma striatum</i> (von Schlotheim); Foster & Sebe, p. 168, fig. 3M.
19		
20		
21		
22	1328	
23		
24		
25	1329	<i>Material.</i> 49 LVs, 30 RVs and nine unassigned shell fragments.
26		
27		
28	1330	
29		
30		
31	1331	<i>Description.</i> Shell medium sized, ovate, higher than long, procrecent, equivalve, inequilateral,
32		
33	1332	moderately inflated; beak prosogyrate, obtuse; dorsal margin angulated, antero-dorsal margin
34		
35	1333	shorter than postero-dorsal one, anterior margin mostly straight, occasionally slightly concave;
36		
37	1334	auricles obtuse and faintly demarcated from flank, anterior auricles smaller than posterior ones,
38		
39	1335	small byssal gape between valves below anterior auricle; lunula large, elongated, impressed, located
40		
41		
42	1336	dorsally to anterior umbonal ridge and comprising nearly the entire antero-dorsal margin (Fig. 6I); up
43		
44	1337	to 60 radial ribs that start close to beak, crests flattened but getting rounded towards ventral margin,
45		
46	1338	very narrow and punctate interspaces getting broader towards ventral margin where they are almost
47		
48	1339	as broad as the ribs, number of ribs increasing by bifurcation, occasionally ribs and interspaces
49		
50	1340	getting broader distally but sometimes also remaining relatively narrow, occasionally ribs curved on
51		
52		
53	1341	ventral parts of shell, auricles bearing ribs and occasionally weak growth lines, growth lines strong on
54		
55	1342	lunula and anterior auricles.
56		
57		
58	1343	
59		
60		

- 1344 *Remarks.* This well-known Middle Triassic species is very abundant in the samples from Dobrogea.
- 1345 Our material agrees very well with specimens from the type area in the Germanic Basin, although
- 1346 some specimens have a slightly higher number of ribs (up to 60) than typical specimens from the
- 1347 Muschelkalk (30–50; Schmidt 1928).
- 1348
- 1349 *Ecology.* Species of the genus *Plagiostoma* are interpreted as epibyssate suspension-feeders
- 1350 (Seilacher 1954; McRoberts 2017).
- 1351
- 1352 *Occurrence.* Gogolin Beds (Poland, late Olenekian – early Pelsonian; Brom *et al.* 2016), Jena
- 1353 Formation (Germany, late Bithynian – early Illyrian; Wagner 1897; Assmann 1937; Schmidt 1928,
- 1354 1938; Klotz & Lukas 1988), Valle Trompia (Italy, middle Anisian; Speciale 1967), Felsőőrs Formation
- 1355 (Hungary, Pelsonian–Illyrian; Szente & Vörös 2003), Zuhány Limestone (Hungary, late Pelsonian –
- 1356 early Illyrian; Foster & Sebe 2017), Leidapo Member (China, late Pelsonian – early Illyrian; Stiller
- 1357 2001; Komatsu *et al.* 2004), Karlstadt Formation, Heilbronn Formation, Diemel Formation (Germany,
- 1358 early Illyrian; Assmann 1937; Schmidt 1928, 1938), Trochitenkalk Formation (Germany, middle
- 1359 Illyrian; Wagner 1897; Schmidt 1928, 1938; Busse 1972; Hagdorn & Ockert 1993; Ockert 1993),
- 1360 Meissner Formation (Germany, late Illyrian – early Longobardian; Wagner 1897; Schmidt 1928, 1938;
- 1361 Busse 1972; Hagdorn & Mundlos 1982; Hagdorn & Ockert 1993), Dont Formation (Italy, Pelsonian–
- 1362 Illyrian; Posenato 2008), Erfurt Formation (Germany, early Longobardian; Schmidt 1928, 1938;
- 1363 Hagdorn 2015).

1364

1365 *Plagiostoma cf. angulatum* (Münster, 1841)

1366 Figures 6M–N

1367

1		
2		
3	1368	cf. *1841 <i>Lima angulata</i> ; Münster, p. 73, pl. 6, fig. 30.
4		
5		
6	1369	cf. 1895 <i>Lima angulata</i> Münster; Bittner, p. 173, pl. 22, figs 3–5.
7		
8		
9	1370	cf. 1903 <i>Lima angulata</i> Münster; Broili, p. 176, pl. 20, figs 1–3, 10.
10		
11		
12	1371	cf. 1909 <i>Lima angulata</i> Münster; Wilckens, p. 132, pl. 5, fig. 15.
13		
14		
15	1372	cf. 1923 <i>Lima angulata</i> Münster; Diener, p. 102 [cum syn.].
16		
17		
18	1373	cf. 1981 <i>Plagiostoma angulata</i> [sic] (Münster); Zardini, pl. 17, fig. 24.
19		
20		
21	1374	
22		
23		
24	1375	<i>Material.</i> One LV and two RVs, one of them an internal mould with some shell remains.
25		
26		
27	1376	
28		
29		
30	1377	<i>Description.</i> Shell medium sized, inequilateral, obliquely ovate in outline with the axis of greatest
31		
32	1378	elongation oriented from postero-dorsal to antero-ventral, higher than long, procrescent,
33		
34	1379	moderately inflated; beak slightly prosogyrate, acute, slightly projecting above short hinge margin;
35		
36	1380	anterior shell margin straight; lunula triangular, elongated, located dorsally of anterior umbonal
37		
38	1381	ridge; auricles small, obtuse and faintly demarcated from flank, anterior auricle smaller than
39		
40	1382	posterior one; radial ribs with rounded crests that start close to beak, up to 25 first order ribs getting
41		
42	1383	broader ventrally, also visible on internal mould, interspaces between first order ribs variable in
43		
44	1384	width, covered with several (up to three) fine radial ribs that are not visible on internal moulds,
45		
46	1385	posterior auricle with radial ribs, anterior auricle with well-developed incremental lines.
47		
48		
49	1386	
50		
51		
52	1387	<i>Remarks.</i> The Dobrogea specimens are morphologically similar to <i>Lima angulata</i> Münster, 1841.
53		
54	1388	However, Bittner (1895) mentioned that the higher order ribs in this species vary in thickness and
55		
56	1389	few can become as broad as the first order ribs. Furthermore, higher order ribs can be located on the
57		
58		
59		
60		

1390 flanks of first order ribs in larger specimens with thin first order ribs. Both features were not  
1391 observed in the Dobrogea specimens. Bitter (1895) further mentioned that higher order ribs are  
1392 more abundant on the posterior part. In the Dobrogea specimens such a dominance of higher order  
1393 ribs in the posterior part of the shell is not observable. Although Bittner (1895) pointed out that  
1394 irregularities concerning the number, strength and distribution of ribs are a conspicuous feature of  
1395 this species, we prefer to retain this species in open nomenclature.

1396

1397 *Ecology.* Same as for *Plagiostoma striatum*.

1398

1399 *Occurrence.* Cima Viezzena (Italy, Longobardian; Wilckens 1909), Pachycardientuffe (Italy, late  
1400 Longobardian; Broili 1903; Blaschke 1905; Waagen 1907), Cassian Formation (Italy, Longobardian–  
1401 Julian; Bittner 1895; Pia 1937; Zardini 1981).

1402

1403 Subclass PALAEOHETERODONTA Newell, 1965

1404 Order TRIGONIIDA Dall, 1889

1405 Superfamily MYOPHORIIDEA Bronn, 1849 in Bronn 1848–1849

1406 Family MYOPHORIIDAE Bronn, 1849 in Bronn 1848–1849

1407 Genus ELEGANTINIA Waagen, 1907

1408

1409 *Type species.* *Lyrodon elegans* Dunker, 1851.

1410

1  
2  
3 1411 *Remarks.* The classification of *Elegantinia* and the closely related genus *Gruenewaldia* within  
4  
5 1412 Myophoriidae, Gruenewaldiidae or Trigoniidae has been debated (e.g. Newell & Boyd 1975, 1987;  
6  
7 1413 Cooper 1991; Boyd & Newell 1997). Based on hinge morphology and the absence of trigonioid pedal  
8  
9 1414 muscles, Hautmann (2003) suggested retaining these genera within Myophoriidae, which is followed  
10  
11  
12 1415 herein.

13  
14  
15 1416

16  
17  
18 1417 *Elegantinia elegans* (Dunker, 1849)

19  
20 1418 Figures 7A–B

21  
22  
23 1419

24  
25  
26 1420 \*1849 *Lyriodon elegans*; Dunker, p. 15.

27  
28  
29 1421 1928 *Myophoria elegans* (Dunker); Schmidt, p. 193, fig. 455.

30  
31  
32 1422 1975 *Lyriomyophoria elegans* (Dunker); Newell & Boyd, p. 148, fig. 89.

33  
34  
35 1423 1976 *Myophoria (Elegantinia) elegans* (Dunker); Gu *et al.*, pl. 21, figs 1–9.

36  
37  
38 1424 2004 *Elegantinia elegans* (Dunker); Komatsu *et al.*, p. 216, figs 12–13.

39  
40  
41 1425 2015 *Elegantinia elegans* (Dunker); Hagdorn, p. 120, fig. 7.20.

42  
43  
44 1426

45  
46  
47 1427 *Material.* One almost complete RV.

48  
49  
50 1428

51  
52  
53 1429 *Description.* Shell medium sized, longer than high, subtrigonal, inequilateral, moderately inflated,

54  
55 1430 anterior and ventral margin rounded, posterior margin truncated; beak not preserved; marginal

56  
57 1431 carina prominent, passing from umbo to postero-ventral margin, becoming broader distally;

58  
59 1432 antecarinal depression present; radial groove on corselet; at least 21 strong, commarginal ribs on

1433 flank, getting narrower toward ventral margin, interspaces almost as broad as ribs, corselet with one  
 1434 radial rib behind radial groove, more and finer commarginal ribs than on flank, no ornamentation  
 1435 visible on marginal carina and antecarinal depression; hinge with two smooth teeth, merged below  
 1436 the beak (Fig. 7B).

1437

1438 *Remarks.* The Dobrogea specimen agrees in all observed characters with the morphology of  
 1439 *Elegantinia elegans*, a species that is widespread in the Muschelkalk of the Germanic Basin but also  
 1440 reported from the Anisian of China.

1441

1442 *Ecology.* Shallow infaunal suspension-feeder.

1443

1444 *Occurrence.* Phra That Fm. (Thailand, Lower Triassic; Vu Khuc 2000), Röt Formation (Germany, early  
 1445 Bithynian; Schmidt 1928, 1938), Jena Formation (Germany, late Bithynian – early Illyrian; Wagner  
 1446 1897; Schmidt 1928, 1938), Leidapo Member (China, late Pelsonian – early Illyrian; Gu *et al.* 1976;  
 1447 Stiller 2001; Komatsu *et al.* 2004), Karlstadt Formation, Heilbronn Formation, Diemel Formation  
 1448 (Germany, early Illyrian; Schmidt 1928, 1938), Trochitenkalk Formation (Germany, middle Illyrian;  
 1449 Wagner 1897; Schmidt 1928, 1938; Busse 1972; Hagdorn & Ockert 1993; Ockert 1993), Meissner  
 1450 Formation (Germany, late Illyrian – early Longobardian; Wagner 1897; Schmidt 1928, 1938; Hagdorn  
 1451 & Ockert 1993; Ockert 1993), Erfurt Formation (Germany, early Longobardian; Schmidt 1928, 1938;  
 1452 Hagdorn 2015).

1453

1454 Genus HEMINAJAS Neumayr, 1891

1455



1		
2		
3	1456	<i>Type species. Myophoria fissidentata</i> von Wöhrmann, 1889.
4		
5		
6	1457	
7		
8		
9	1458	<i>Remarks.</i> We follow Hautmann <i>et al.</i> (2013) and place <i>Heminajas</i> , which has three teeth in its RV,
10		
11	1459	within the Myophoriidae, in which a relatively weak third tooth may be present in their RVs, and not
12		
13		
14	1460	in the Trigonodidae (= Pachycardiidae in Cox <i>et al.</i> 1969), which do not possess more than two teeth
15		
16	1461	in each of their valves (Cox <i>et al.</i> 1969).
17		
18		
19	1462	
20		
21		
22	1463	<i>Heminajas?</i> sp.
23		
24	1464	Figure 7C
25		
26		
27	1465	
28		
29		
30	1466	<i>Material.</i> One almost completely preserved RV.
31		
32		
33	1467	
34		
35		
36	1468	<i>Description.</i> Shell medium sized, elliptical, longer than high, inequilateral, retrocrescent; beak
37		
38		
39	1469	positioned in anterior part of shell, projecting dorsal margin; slight depression in front of the umbo
40		
41	1470	creating a flattening of the shell towards the anterior margin; regular growth lines, regular radial
42		
43	1471	lines best visible at anterior and posterior part of the shell, growth lines and radial lines create a
44		
45	1472	weak reticular pattern.
46		
47		
48		
49	1473	
50		
51		
52	1474	<i>Remarks.</i> The morphology of the Dobrogea specimen is similar to that of <i>Heminajas</i> . However,
53		
54	1475	neither a weak umbonal carina nor the truncated posterior margin are present/preserved in the
55		
56	1476	Dobrogea specimen. Furthermore, the internal characters are not observable, which makes the
57		
58	1477	generic assignment uncertain. The radial lines may result from the shell microstructure.
59		
60		

1  
2  
3 1478  
4  
5

6 1479 *Ecology.* Shallow infaunal suspension-feeder.  
7  
8

9 1480  
10  
11

12 1481 Order MODIOMORPHOIDA Cox *et al.*, 1969  
13  
14

15 1482 Superfamily MODIOMORPHOIDEA Miller, 1877  
16  
17

18 1483 Family HEALEYIDAE Hautmann, 2008  
19  
20

21 1484 Genus JOANNINA Waagen, 1906  
22  
23

24 1485  
25

26  
27 1486 *Type species.* *Joannina joannae* Waagen, 1906.  
28  
29

30 1487  
31  
32

33 1488 *Remarks.* We follow Hautmann (2008) and place *Joannina* within Healeyidae and this family within  
34

35 1489 Modiomorphoidea.  
36  
37

38 1490  
39  
40

41 1491 *Joannina* aff. *joannae* Waagen, 1906  
42  
43

44 1492 Figure 7D  
45  
46

47 1493  
48  
49

50 1494 aff. \*1906 *Joannina joannae*; Waagen, p. 390.  
51  
52

53 1495 aff. 1907 *Joannina joannae* Waagen; Waagen, p. 34, pl. 34, figs 12–14.  
54  
55

56 1496  
57  
58

59 1497 *Material.* One RV.  
60

1		
2		
3	1498	
4		
5		
6	1499	<i>Description.</i> Shell medium sized, subrectangular-modioliform, retrocrescent; beak subterminal and
7		
8	1500	prosogyrate, giving rise to pronounced diagonal ridge; diagonal ridge sharp near beak, becoming
9		
10	1501	flatter and broader towards postero-ventral margin, separating flank from posterior area; median
11		
12	1502	depression on center of flank separating anterior shell lobe; posterior area with blunt, faint radial
13		
14	1503	structure but no distinct ridges; ornamentation with several pronounced growth irregularities and
15		
16	1504	fine incremental lines becoming irregular to weavy on posterior area.
17		
18		
19		
20	1505	
21		
22		
23	1506	<i>Remarks.</i> This species is similar to <i>Joannina joannae</i> Waagen, 1906 from the late Ladinian, but it
24		
25	1507	differs in being more rectangular in outline and lacking two sharp ridges on the posterior area (see
26		
27	1508	Hautmann 2008 for a description of <i>J. joannae</i> ). These differences are probably sufficient for
28		
29	1509	distinguishing this taxon as a new species, but with the limited material available to us, we
30		
31	1510	provisionally describe it in open nomenclature.
32		
33		
34		
35	1511	
36		
37		
38	1512	<i>Ecology.</i> Hautmann (2008) suggested an endobyssate mode of life for <i>Joannina</i> because of its
39		
40	1513	modioliform shell and the presence of a byssal gape.
41		
42		
43	1514	
44		
45		
46		
47	1515	Family KALENTERIDAE Marwick, 1953
48		
49		
50	1516	Genus MYOCONCHA J. Sowerby, 1824
51		
52	1517	
53		
54		
55	1518	<i>Type species.</i> <i>Myoconcha crassa</i> J. Sowerby, 1824.
56		
57		
58	1519	
59		
60		

1520 *Remarks.* The 'actinodont' hinge of the Kalenteridae indicates their affiliation to the  
 1521 Modiomorphoidea and Palaeoheterodonta (Hautmann 2001a). In contrast, Kaim & Schneider (2012)  
 1522 placed the Kalenteridae in the Carditida within Heteroconcha, because the inner shell layer of  
 1523 *Myoconcha* and *Caspiconcha* is cross-lamellar, rather than nacreous. However, the crossed-lamellar  
 1524 shell structure developed several times independently, e.g. in Pectinida and Heterodonta (e.g. Carter  
 1525 1990). We therefore regard the hinge morphology as a more important taxonomic character than the  
 1526 shell structure.

1528 *Myoconcha* sp.

1529 Figure 7E

1531 *Material.* One posterior fragment of a RV.

1533 *Description.* Medium sized fragment of posterior part of shell externally covered by at least 12  
 1534 narrow radial ribs and straight, sub-commarginal rib-like growth lines, forming tubercles where  
 1535 intersecting radial ribs.

1537 *Remarks.* The shape and ornamentation of the studied fragment are indicative of *Myoconcha*. The  
 1538 sculpture and number of ribs of the described specimen show most similarities with *Myoconcha*  
 1539 *appeli* Kittl, 1903, *Myoconcha maximiliani-leuchtenbergensis* von Klipstein, 1843 and *Myoconcha*  
 1540 *pavida* Bittner, 1901, but incomplete preservation prevents us from a firm assignment.

1541

1		
2		
3	1542	<i>Ecology</i> . Hodges (2000) suggested an endobyssate mode of life for oval-modioliform specimens of
4		
5	1543	<i>Myoconcha</i> , whereas Kaim & Schneider (2012) suggested an epifaunal mode of life for mytiliform
6		
7	1544	<i>Myoconcha</i> specimens. Because of its incomplete preservation the exact mode of life for the
8		
9		
10	1545	Dobrogea specimen remains unclear.
11		
12		
13	1546	
14		
15		
16	1547	Subclass HETERODONTA Neumayr, 1884
17		
18		
19	1548	Infraclass EUHETERODONTA Giribet & Distel, 2003
20		
21		
22	1549	Order LUCINIDA Gray, 1854
23		
24		
25	1550	Superfamily LUCINOIDEA Fleming, 1828
26		
27		
28	1551	Family MACTROMYIDAE Cox, 1929
29		
30		
31	1552	Genus UNICARDIUM d’Orbigny, 1849
32		
33		
34	1553	
35		
36		
37	1554	<i>Type species</i> . <i>Corbula cardioides</i> Phillips, 1829.
38		
39		
40	1555	
41		
42		
43	1556	<i>Unicardium schmidii</i> (Geinitz, 1842)
44		
45	1557	Figures 7F–H
46		
47		
48	1558	
49		
50		
51	1559	*1842 <i>Arca? schmidii</i> ; Geinitz, p. 557, pl. 10, fig. 9.
52		
53		
54	1560	1859 <i>Arca? schmidii</i> Geinitz; von Schauroth, p. 323, pl. 2, fig. 16.
55		
56		
57	1561	1923 <i>Unicardium schmidi</i> (Geinitz); Diener, p. 121 [cum syn.].
58		
59		
60	1562	1928 <i>Unicardium schmidi</i> (Geinitz); Schmidt, p. 199, fig. 476.

1563

1564 *Material.* Six LVs, three RVs and three additional specimens, mostly incompletely preserved. Two of  
1565 them are preserved as internal moulds.

1566

1567 *Description.* Shell medium sized, elliptical, equivalve, slightly inequilateral, moderately inflated; beak  
1568 strongly prosogyrate, obtuse, located almost in the middle of straight dorsal margin and distinctly  
1569 projecting above it; ligament parivincular, nymph long and straight (Fig. 7G); hinge edentulous;  
1570 postero-dorsal margin slightly higher than antero-dorsal margin; strong commarginal ribs with  
1571 interspaces becoming narrower towards ventral margin, growth lines visible in interspaces.

1572

1573 *Remarks.* The Dobrogea specimens agree very well with *Unicardium schmidii*, as described in Schmidt  
1574 (1928), including the edentulous hinge. However, *Unicardium* d'Orbigny, 1849 was originally  
1575 described as having a tooth in its hinge (d'Orbigny 1849, p. 218), and the question arises whether the  
1576 edentulous *U. schmidii* really belongs to this genus or, more generally, whether the presence or  
1577 absence of a hinge tooth is within the variability of the genus. Unfortunately, the hinge of the type  
1578 species of *Unicardium*, *Corbula cardioides* Phillips, 1829, was neither described nor illustrated so far.  
1579 Stoliczka (1871, p. 246) noted that *Unicardium* has a hinge with one cardinal tooth in each valve that  
1580 is occasionally obsolete or nearly so, which seems to justify the assignment of *U. schmidii* to this  
1581 genus.

1582

1583 *Ecology.* Moderately deep infaunal suspension-feeder.

1584

1		
2		
3	1585	<i>Occurrence.</i> Röt Formation (Germany, early Bithynian; Schmidt 1928, 1938), Jena Formation
4		
5	1586	(Germany, late Bithynian – early Illyrian; Wagner 1897; Schmidt 1928, 1938; Klotz & Lukas 1988),
6		
7	1587	Trochitenkalk Formation (Germany, middle Illyrian; Wagner 1897; Schmidt 1928, 1938; Ockert 1993),
8		
9		
10	1588	Meissner Formation (Germany, late Illyrian – early Longobardian; Wagner 1897; Schmidt 1928, 1938;
11		
12	1589	Busse 1972; Hagdorn und Ockert 1993), Erfurt Formation (Germany, early Longobardian; Schmidt
13		
14	1590	1928, 1938).
15		
16		
17	1591	
18		
19		
20	1592	Family LUCINIDAE Fleming, 1828
21		
22		
23	1593	Genus SCHAFHAEUTLIA Cossmann, 1897
24		
25		
26	1594	
27		
28		
29	1595	<i>Type species.</i> <i>Gonodon schafhaeutli</i> Salomon, 1895.
30		
31		
32	1596	
33		
34		
35	1597	<i>Remarks.</i> We follow the conventional assignment of <i>Schafhaeutlia</i> to the Lucinidae (e.g. Cox <i>et al.</i>
36		
37	1598	1969), but we note that the morphological evidence for this is questionable.
38		
39		
40	1599	
41		
42		
43	1600	<i>Schafhaeutlia astartiformis</i> (Münster, 1841)
44		
45	1601	Figures 7I–K
46		
47		
48	1602	
49		
50		
51		
52	1603	*1841 <i>Isocardia astartiformis</i> ; Münster, p. 87, pl. 8, fig. 24.
53		
54		
55	1604	1865 <i>Corbis astartiformis</i> (Münster); Laube, p. 37, pl. 15, fig. 5.
56		
57		
58	1605	1895 <i>Gonodon astartiformis</i> (Münster); Bittner, p. 12, pl. 3, figs 1–4.
59		
60		
	1606	1903 <i>Gonodon astartiformis</i> (Münster); Broili, p. 218, pl. 27, figs 30–31.

1607 1909 *Schafhäutlia astartiformis* (Münster); Wilckens, p. 162, pl. 6, fig. 11.

1608 1923 *Schafhäutlia astartiformis* (Münster); Diener, p. 217 [cum syn.].

1609 1966 *Gonodon astartiformis* (Münster); Allasinaz, p. 701, pl. 54, figs 6–7.

1610 1981 *Schafhaeutlia astartiformis* (Münster); Zardini, pl. 33, figs 8a–b; pl. 34, figs 1–6.

1611

1612 *Material.* Four LVs, one RV and one articulated specimen. Mostly incompletely preserved or internal  
1613 moulds.

1614

1615 *Description.* Shell medium sized, moderately inflated, slightly inequilateral, elliptical, longer than  
1616 high; beak strongly prosogyrate, obtuse, located close to the middle of dorsal margin and clearly  
1617 projecting above it; nymph long and straight; ligament parivincular; cardinal tooth 3a observed in  
1618 one specimen (Fig. 7J); irregular growth lines variable in strength, occasionally forming weak ribs.

1619

1620 *Remarks.* The overall shape in combination with the morphology of cardinal tooth 3a (Fig. 7J) is  
1621 indicative of *Schafhaeutlia*. However, already Bittner (1901) noted that species of this genus are  
1622 difficult to distinguish from each other. We found the best agreement with *Schafhaeutlia*  
1623 *astartiformis*, a well-known species that occurs in many Middle and Late Triassic localities.

1624

1625 *Ecology.* Hautmann (2001a) proposed an epibenthic mode of life for *Schafhaeutlia sphaerioides*.  
1626 According to Hautmann (2001a), *S. sphaerioides* rested on its flattened anterior side with the  
1627 strongly curved beaks stabilizing this position. However, the anterior margin of *S. astartiformis* is well  
1628 rounded and not flattened, suggesting a shallow infaunal mode of life.



1		
2		
3	1629	
4		
5		
6	1630	<i>Occurrence.</i> Leidapo Member (China, late Pelsonian – early Illyrian; Stiller 2001; Komatsu <i>et al.</i> 2004),
7		
8	1631	Cima Viezzena (Italy, Longobardian; Wilckens 1909), Pachycardientuffe (Italy, late Longobardian;
9		
10	1632	Broili 1903; Blaschke 1905), Cassian Formation (Italy, Longobardian–Julian; Laube 1865; Bittner 1895;
11		
12	1633	Fürsich & Wendt 1977; Zardini 1981), Sella delle Cave (Italy, Julian; Allasinaz 1966).
13		
14		
15		
16	1634	
17		
18		
19	1635	<i>Schafhaeutlia? trigona</i> (Stoppani, 1858)
20		
21	1636	Figure 7L
22		
23		
24	1637	
25		
26		
27	1638	*1858 <i>Cyprina trigona</i> ; Stoppani, p. 85, pl. 17, fig. 8.
28		
29		
30	1639	
31		
32		
33	1640	<i>Material.</i> Two LVs.
34		
35		
36	1641	
37		
38		
39	1642	<i>Description.</i> Shell small to medium sized, elliptical, longer than high, slightly inflated, inequilateral;
40		
41	1643	beak strongly prosogyrate, obtuse, clearly projecting above straight dorsal margin, positioned at the
42		
43	1644	anterior part of the shell; fine, regular growth lines well visible with naked eye.
44		
45		
46		
47	1645	
48		
49		
50	1646	<i>Remarks.</i> The main features of the Dobrogea specimens are in accordance with the main
51		
52	1647	characteristics of <i>Cyprina trigona</i> . Rossi Ronchetti (1959) assigned a morphologically similar species
53		
54	1648	to <i>Schafhaeutlia</i> . Although the Dobrogea specimens, as well as the specimens illustrated in Stoppani
55		
56	1649	(1858) and Rossi Ronchetti (1959), resemble the shape of <i>Schafhaeutlia</i> , the anteriorly positioned
57		
58	1650	beak is unusual for this genus. Unfortunately, the internal characters of the Dobrogea specimens are
59		
60		

- 1651 not observable, which makes an assignment at the genus level uncertain. This species differs from
- 1652 *Schafhaeutlia astartiformis* by its shape and the fine and regular growth lines.
- 1653
- 1654 *Ecology*. Same as for *Schafhaeutlia astartiformis*.
- 1655
- 1656 *Occurrence*. Esino Limestone (Italy, Ladinian; Stoppani 1858).
- 1657
- 1658 Order MEGALODONTIDA Starobogatov, 1992
- 1659 Superfamily MEGALODONTOIDEA Morris & Lycett, 1853
- 1660 Family DICEROCARDIIDAE Kutassy, 1934?
- 1661 Genus PRAEDICEROCARDIUM n. gen.
- 1662
- 1663 *LSID*. urn:lsid:zoobank.org:act:A36FA7B2-1F30-497E-B56C-C791F54E1873
- 1664
- 1665 *Derivation of name*. Combination of *prae* (Latin: before) and the genus name *Dicerocardium*.
- 1666
- 1667 *Type species*. *Praedicerocardium vetulus* n. gen. et n. sp.
- 1668
- 1669 *Diagnosis*. Shell equivalve; valves triangular; flank delimited anteriorly and posteriorly by sharp ridges
- 1670 converging towards the umbo; angle included between ridges acute; shell region in front of anterior

1  
2  
3 1671 ridge flattened; shallow but broad radial shell depressions in front of posterior ridge and narrow  
4  
5 1672 escutcheon behind it; beaks terminal; umbones strongly prosogyrate but not elevated above flat  
6  
7 1673 anterior shell region.  
8  
9  
10  
11 1674  
12  
13  
14 1675 *Remarks:* Because the hinge structure is not observable, the external shell morphology is the sole  
15  
16 1676 basis for a discussion about the taxonomic position of the genus. The new genus closely resembles  
17  
18 1677 species assigned to *Dicerocardium* Stoppani, 1865 (e.g. *Dicerocardium dolomiticum marianii* Frech,  
19  
20 1678 1904) in terms of its triangular shape, flattened anterior shell region and the presence of an anterior  
21  
22  
23 1679 and posterior carina. However, it differs from *Dicerocardium* by the morphology of the umbones,  
24  
25 1680 which are less incoiled and not elevated above the flat anterior shell region. Moreover, the  
26  
27 1681 stratigraphical occurrences differ considerably: *Dicerocardium* is Norian (Végh-Neubrandt 1982), thus  
28  
29 1682 appearing about 20 million years later than *Praedicerocardium*.  
30  
31  
32  
33 1683 Based on the similarity in the external morphology, we place the new genus tentatively in  
34  
35 1684 Dicerocardiidae, rather than in Megalodontidae. If correct, the new genus predates considerably the  
36  
37 1685 oldest stratigraphic appearance of the Dicerocardiidae from the Carnian (Végh-Neubrandt 1982, Tab.  
38  
39 1686 18) to the Anisian. Moreover, it also predates the stratigraphical oldest Triassic megalodontid, which  
40  
41 1687 is a (questionable) report from the Meissner Formation (*'Megalodon securiformis'* Schmidt, 1928)  
42  
43 1688 that is late Illyrian (late Anisian) to early Longobardian (late Ladinian) in age (Hagdorn & Nitsch 2009).  
44  
45  
46 1689 If Dicerocardiidae evolved from Megalodontidae (Cox *et al.* 1969; Végh-Neubrandt 1982), the early  
47  
48 1690 stratigraphic appearance of Dicerocardiidae in the Middle Triassic implies a considerable outage in  
49  
50 1691 the fossil record of Megalodontidae during the Early and early Middle Triassic.  
51  
52  
53 1692 The new genus has also implications for the phylogeny of Dicerocardiidae. Végh-Neubrandt (1982, p.  
54  
55 1693 116–117) suggested that *Dicerocardium* derives from a carnian lineage that led from *Laubeia* via  
56  
57  
58 1694 *Physocardia* and *Cornucardia* to the Norian *Dicerocardium*. We suggest that *Praedicerocardium* is  
59  
60 1695 another candidate ancestor, which is in its external morphology much closer to *Dicerocardium* than is

1696 *Physocardia*. Future data on the hinge structure may help to decide whether this similarity in the  
 1697 external morphology resulted from phylogenetic relationships of convergent evolution.

1698

1699 *Praedicerocardium vetulus* n. gen. et n. sp.

Figures 7M–P

1701

1702 *LSID*. urn:lsid:zoobank.org:act:F1CC713D-556D-4971-AA99-B0253CA28F79

1703

1704 *Derivation of name*. *Vetulus* (Latin: old), because it is the geologically oldest Triassic megalodontoid.

1705

1706 *Holotype*. LPB IIL 2142

1707 *Paratypes*. LPB IIL 2143, IIL 2144

1708

1709 *Material*. One LV, two RVs and one articulated specimen.

1710

1711 *Diagnosis*: Same as for the genus.

1712

1713 *Description*. Shell large, trigonal, longer than high, highly inequilateral, strongly inflated; shell thick,  
 1714 consisting of two shell layers, the outer layer approximately 1 mm thick, the inner layer  
 1715 approximately 3–4 mm thick; beaks strongly coiled, prosogyrate, terminal, clearly projecting above  
 1716 dorsal margin; anterior part of shell flat, separated from flank by sharp ridge extending from umbo to

1  
2  
3 1717 antero-ventral margin; escutcheon narrow, separated from flank by sharp escutcheon ridge, shallow  
4  
5 1718 but broad radial shell depression anterior to escutcheon ridge; anterior margin straight, posterior  
6  
7 1719 margin slightly curved, distally almost straight; regular growth lines, fine and slightly irregular  
8  
9  
10 1720 commarginal ribs present in one specimen; internal structures not observed.  
11  
12  
13 1721  
14  
15  
16 1722 *Ecology*. According to Zapfe (1957) and Végh-Neubrandt (1982), equivalved megalodontids were  
17  
18 1723 semi-infaunal or epifaunal suspension-feeders that rested on their anterior shell margin and beaks. In  
19  
20 1724 this position, the flat and relatively large anterior part of *Praedicerocardium vetulus* provided a  
21  
22 1725 resting surface and prevented the animal from sinking into the soft sediment. Alternatively,  
23  
24 1726 *Praedicerocardium vetulus* might have rested on the straight postero-dorsal margin, as suggested by  
25  
26 1727 Schneider (2017) for the Jurassic *Pachyrisma*. This position would also be stable, supported by the  
27  
28 1728 escutcheon ridges and the center of gravity of the shell. Furthermore, it would expose the respiratory  
29  
30 1729 margin directly into the free water column. However, it raises the problem how the animal could  
31  
32 1730 adopt this position, given that its foot would also be directed towards the free water column,  
33  
34 1731 without contact to the sediment and therefore essentially inoperable.  
35  
36  
37  
38  
39 1732  
40  
41  
42 1733 Order ANOMALODESMATA Dall, 1889  
43  
44  
45 1734 Suborder PHOLADOMYIDA Newell, 1965  
46  
47  
48 1735 Superfamily PHOLADOMYOIDEA King, 1844  
49  
50  
51 1736 Family MEGADESMIDAE Vokes, 1967  
52  
53  
54 1737 Genus ROMANIAMYA n. gen.  
55  
56  
57 1738  
58  
59  
60 1739 *LSID*. urn:lsid:zoobank.org:act:5CAB38FD-6B23-43C0-A750-2C2E55F501AD

1740

1741 *Derivation of name.* Combination of 'Romania' and the genus name *Mya*.

1742

1743 *Type species.* *Romaniamya mahmudiaensis* n. gen. et n. sp.

1744

1745 *Diagnosis.* Putative megadesmid bivalves with shell slightly expanding posteriorly, having a shallow  
 1746 median depression on the flank, a moderate siphonal gape, a straight dorsal margin with short  
 1747 anterior and posterior projections and a blunt posterior umbonal ridge.

1748

1749 *Remarks.* This enigmatic taxon shows some similarities with late Palaeozoic anomalodesmats that are  
 1750 usually referred to the family Megadesmidae (e.g. Cox *et al.* 1969; Runnegar 1974), most notably  
 1751 with the Permian *Pachymyonia* Dun, 1932. Some of these megadesmids possess a well-developed  
 1752 umbonal ridge, short distal projections of the nearly straight dorsal margin, a shallow depression on  
 1753 the flank and a siphonal gape, but a combination of these characters as in the Dobrogea specimens is  
 1754 unknown so far.

1755

1756 *Romaniamya mahmudiaensis* n. gen. et n. sp.

1757 Figures 8A–B

1758

1759 *LSID.* urn:lsid:zoobank.org:act:4E73DC0A-5A9D-4D76-9943-3339A2A49CB7

1760

1761 *Derivation of name.* Referring to the type locality near Mahmudia, North Dobrogea, Romania.

1		
2		
3	1762	
4		
5		
6	1763	<i>Holotype</i> . LPB IIL 2145
7		
8		
9	1764	<i>Paratype</i> . LPB IIL 2146
10		
11		
12	1765	
13		
14		
15	1766	<i>Material</i> . One LV and internal moulds of an articulated specimen and a LV.
16		
17		
18	1767	
19		
20		
21	1768	<i>Diagnosis</i> . Same as for the genus.
22		
23		
24	1769	
25		
26		
27	1770	<i>Description</i> . Shell medium sized, elongated, trapeziform, inequilateral, equivalve, moderately
28		
29	1771	inflated; very shallow median depression on centre of flank; beaks slightly prosogyrate, obtuse,
30		
31	1772	positioned in front of the midline of straight dorsal margin and slightly projecting above it; umbonal
32		
33	1773	ridge prominent, broad, starting close to beak and running towards the postero-ventral part of the
34		
35	1774	shell; sulcus located dorsally to umbonal ridge (Fig. 8B); postero-dorsal margin straight, ventral
36		
37	1775	margin almost straight; slightly gaping at posterior end (Fig. 8B); irregular and weak commarginal ribs
38		
39	1776	present; internal structures unknown.
40		
41		
42		
43	1777	
44		
45		
46	1778	<i>Remarks</i> . <i>Macrodon taramellii</i> Parona, 1889 is very similar to <i>Romaniamya mahmudiaensis</i> but it
47		
48	1779	differs from <i>R. mahmudiaensis</i> by the shape of the antero-dorsal margin and the presence of fine
49		
50	1780	radial striae. Parona (1889) noted that the ligament area was not observable in his specimens,
51		
52	1781	therefore his assignment to <i>Macrodon</i> is uncertain.
53		
54		
55	1782	
56		
57		
58		
59		
60		

1783 *Ecology.* The posterior gape suggests the presence of well-developed siphons and therefore a  
1784 deep-infaunal mode of life.

1785

1786 Superfamily UNCERTAIN

1787 Family UNCERTAIN

1788 Genus UNCERTAIN

1789

1790 Anomalodesmata indet. A

1791 Figure 8C

1792

1793 *Material.* One incompletely and poorly preserved internal mould of a RV.

1794

1795 *Description.* Shell medium sized, elongated, slightly inflated, strongly inequilateral, retrocrescent;  
1796 beak close to anterior margin, slightly projecting long dorsal margin; dorsal and ventral margin  
1797 subparallel to each other.

1798

1799 *Remarks.* The shape of this internal mould suggests an affiliation to the Anomalodesmata, but it is  
1800 too incompletely preserved for a more precise determination.

1801

1802 *Ecology.* Deep infaunal suspension-feeder.

1803



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1804

Anomalodesmata indet. B
- 1805

Figure 8D
- 1806
- 1807

*Material.* One fragmented internal mould, possibly representing the posterior part of a RV.
- 1808
- 1809

*Description.* Fragment medium sized, elongated, slightly inflated, becoming broader towards
- 1810

posterior margin; irregular commarginal folds best visible at the margin.
- 1811
- 1812

*Remarks.* In spite of its poor and incomplete preservation, the shape of this fragment is suggestive
- 1813

for an anomalodesmatid. The preservation state of this specimen does not allow a more accurate
- 1814

assignment.
- 1815
- 1816

*Ecology.* Deep infaunal suspension-feeder.
- 1817
- 1818

Anomalodesmata indet. C
- 1819

Figure 8E
- 1820
- 1821

*Material.* One incompletely preserved internal mould of a LV.
- 1822
- 1823

*Description.* Shell medium sized to large, elongated, moderately inflated, strongly inequilateral,
- 1824

retrocrscent; umbonal region curved; beak broad, ophistogyrate, slightly projecting above very long

1825 and straight dorsal margin, positioned in anterior part of shell; ventral margin slightly concave,  
 1826 posterior margin straight; umbonal ridge passing from umbo towards postero-ventral edge.

1827

1828 *Remarks.* The general shape suggests an assignment to the Anomalodesmata but the incomplete  
 1829 preservation state of the Dobrogea specimen does not allow a definite assignment.

1830

1831 *Ecology.* Deep infaunal suspension-feeder.

1832

1833 INCERTAE SEDIS

1834 Genus MYTILICONCHA Tommasi, 1911

1835

1836 *Type species.* *Mytiliconcha orobica* Tommasi, 1911.

1837

1838 *Remarks.* There has been disagreement concerning the validity of *Mytiliconcha*. Cox (1964) proposed  
 1839 the name *Tommasina* as a replacement name for *Mytiliconcha* Tommasi, 1911, which he thought  
 1840 was preoccupied by *Mytiloconcha* Conrad, 1862. However, because the spelling of Conrad's (1862)  
 1841 *Mytiloconcha* differs in one letter from *Mytiliconcha*, the introduction of a replacement name by Cox  
 1842 (1964) was unnecessary (e.g. Vokes 1980; Stiller & Chen 2006; Ros-Franch *et al.* 2014). Waller &  
 1843 Stanley (2005) and Ros-Franch *et al.* (2014) regarded *Mytiliconcha* as a synonym of *Protopis*,  
 1844 although the hinge structure of *Mytiliconcha* is unknown. In contrast, Hautmann (2008) noted that  
 1845 the external features of *Mytiliconcha* and *Protopis* differ profoundly. However, other characters of  
 1846 *Mytiliconcha* are still unknown, making the taxonomic position of *Mytiliconcha* uncertain.

1		
2		
3	1847	
4		
5		
6	1848	<i>Mytiliconcha</i> cf. <i>orobica</i> Tommasi, 1911
7		
8	1849	Figures 8F–H
9		
10		
11	1850	
12		
13		
14	1851	cf. *1911 <i>Mytiliconcha orobica</i> ; Tommasi, p. 35, pl. 3, figs 21–23.
15		
16		
17	1852	cf. 1969 <i>Tommasina orobica</i> (Tommasi); Cox <i>et al.</i> , p. N281, fig. C22(2).
18		
19		
20	1853	
21		
22		
23		
24	1854	<i>Material</i> . Three LVs and three RVs.
25		
26		
27	1855	
28		
29		
30	1856	<i>Description</i> . Shell medium sized, retrocrescent, strongly inflated; beak prosogyrate, acute, projecting
31		
32	1857	above dorsal margin; carina sharp, extending from beak to postero-ventral part of the shell,
33		
34	1858	vanishing distally; shell smooth except for very fine and regular growth lines and faint radial lines that
35		
36		
37	1859	are occasionally visible on some specimens.
38		
39		
40	1860	
41		
42		
43	1861	<i>Remarks</i> . The specimens described above agree well in their external morphology with the type
44		
45	1862	material of <i>Mytiliconcha orobica</i> Tommasi, 1911, but this species has been based on rather
46		
47	1863	incomplete material, which makes any assignment tentative.
48		
49		
50	1864	
51		
52		
53	1865	<i>Ecology</i> . Because of the lack of a byssal gape and the strongly inflated valves the Dobrogea
54		
55		
56	1866	specimens might have been epifaunal or semi-infaunal recliners.
57		
58		
59	1867	
60		

1868 *Occurrence*. Ghegna (Italy, Ladinian; Tommasi 1911).

1869

1870

## 1871 **DISCUSSION AND RESULTS**

1872

1873 With a total of 51 species, the Bivalvia is the most diverse invertebrate group of the  
 1874 *Tubiphytes*-microbial buildup in North Dobrogea. These 51 bivalve species belong to 39 genera, 29  
 1875 families, 20 superfamilies, nine orders and three subclasses, respectively (Fig. 9). Based on the  
 1876 presence of closed ostracod carapaces, Forel & Grădinaru (2018) concluded that the fauna is  
 1877 essentially preserved *in situ*. This conclusion is confirmed by the presence of several bivalve shells  
 1878 with conjoined valves. The quality of the preservation of the bivalve fauna is variable. Some  
 1879 specimens are extremely well preserved and show taxonomically significant characters in detail,  
 1880 whereas other species are represented by fragmentary or poorly preserved specimens only. The  
 1881 latter is partly due to the difficult mechanical extraction of the specimens from the hard rock, which  
 1882 possibly also causes a bias towards a lower bivalve richness (see below). Due to incomplete  
 1883 preservation or the lack of information on taxonomically important shell parts, e.g. the hinge and the  
 1884 ligament, 29 taxa are presented in open nomenclature (Fig. 10). On the other hand, seven species  
 1885 and two genera are newly described (Fig. 10). This relatively high number of new species and genera  
 1886 probably indicates a relatively high degree of endemism, which could be related to the unique  
 1887 environment provided by the *Tubiphytes*-microbial buildup.

1888 Most genera and many of the species are already known from several other Middle Triassic localities  
 1889 (Fig. 11). However, only two species, i.e. *Leptochondria viezzenensis* and *Elegantinia elegans* have  
 1890 been reported from the Early Triassic. Six species from the *Tubiphytes*-microbial buildup are known  
 1891 from the Muschelkalk, 13 species from the Western Tethys and eight species from the Eastern

1892 Tethys. Only one species (*Elegantinia elegans*) has also been reported from Northwestern

1893 Panthalassa (Hayami 1975). Five species (*Gervillaria subelegans*, *Leptochondria separata*, *Entolioides*

1894 *deeckeii*, *Mysidioptera fornicata* and *Schafhaeutlia astartiformis*) and four genera (*Atrina*, *Cultriopsis*,

1895 *Gervillaria*, *Mytiliconcha*) found in the Bithynian *Tubiphytes*-Limestone Member have their oldest

1896 occurrence there (Fig. 10). However, these four genera are generally rare, which makes any

1897 statement about their first occurrences uncertain. In summary, the biogeographic character of the

1898 fauna is largely determined by the palaeogeographic distance to contemporaneous occurrences, plus

1899 an endemic component that might be related to the particular environment.

1900 Ecologically, the bivalve assemblage is clearly dominated by *Plagiostoma striatum* (Fig. 12). Far less

1901 abundant are *Neomorphotis comta*, *Leptochondria pervulgata*, *Chlamys* (*Praechlamys*) *prima*,

1902 *Mysidioptera fornicata* and *Scythentolium anisicum*. All remaining species are relatively rare, and a

1903 total number of 25 species is only represented by one individual. Simpson's index of dominance is

1904 low (0.09) and the trophic nucleus, which is the number of species comprising 80% of the individuals

1905 of an association (Neyman 1967), contains 18 species (Fig. 12). The high evenness could indicate a

1906 heterogeneous environment, a K-selected equilibrium fauna, the effects of time-averaging or a

1907 combination of these factors. The bivalve assemblage contains shallow, moderately deep and deep

1908 infaunal, byssally attached, reclining epifaunal or partly embedded semi-infaunal as well as

1909 cementing species. However, byssal attachment, epifaunal as well as semi-infaunal, represents the

1910 most abundant mode of life, which is shown by the guild-species diversity as well as the absolute

1911 abundance of guilds (Fig. 12). The epibyssate bivalves settled on the hard substratum provided by the

1912 *Tubiphytes*-microbial buildup, where the micro-encruster framework was stabilized by large volumes

1913 of synsedimentary cements (Popa *et al.* 2014). The burrowing, semi-infaunal and free-lying bivalves

1914 indicate the presence of soft substratum, which probably filled pockets between single

1915 *Tubiphytes*-microbial buildups (Fig. 13).

1916 Concerning the richness of bivalve assemblages, a comparison with diversities of other Early and

1917 Middle Triassic bivalve faunas shows that the assemblage from North Dobrogea is more diverse than

1918 all Early Triassic bivalve faunas (Fig. 14 and Table 1). The rarefaction analysis suggests that the bivalve  
1919 richness of the investigated *Tubiphytes*-microbial buildup is undersampled (Fig. 15). Figure 14 and  
1920 Table 1 also show that a set of Middle Triassic bivalve assemblages, including the Jena Formation, the  
1921 Leidapo Member, the Trochitenkalk Formation, the Meissner Formation and the Pachycardientuffe  
1922 are richer than all other Middle Triassic bivalve assemblages. However, because the data used for  
1923 this comparison are raw data from the literature, their interpretation is difficult. Factors like  
1924 oversplitting (see also Friesenbichler *et al.* 2019), a long history of research and a large outcrop area  
1925 (concerning e.g. the Germanic Triassic) or a Lagerstätten effect (e.g. in the Leidapo Member) may  
1926 cause a bias towards higher diversities.

1927 In addition to the 51 bivalves species identified in this study, 21 gastropod species (Nützel *et al.* 2018)  
1928 and 12 brachiopod species (Grădinaru & Gaetani 2019) have been reported so far from this locality.  
1929 These 84 benthic macroinvertebrate species flourished in an environment that was constructed by  
1930 micro-encrusting organisms (Popa *et al.* 2014). Based on the presence of a highly diverse  
1931 invertebrate fauna associated with this microbial buildup, Nützel *et al.* (2018) concluded that benthic  
1932 faunas recovered faster from the end-Permian mass extinction than metazoan reefs, an assumption  
1933 that is in accordance with the results from this study. Similarly, Friesenbichler *et al.* (2019)  
1934 hypothesized that the increase in environmental diversity associated with newly arising carbonate  
1935 factories contributed at least in part to the accelerated pace of recovery during the Middle Triassic.  
1936 The rich Bithynian invertebrate fauna of North Dobrogea associated with one of the earliest large  
1937 Triassic carbonate buildups confirms to this assumption.

## CONCLUSIONS

1  
2  
3 1942 The investigated *in situ* Bithynian (early Middle Triassic) bivalve fauna from the *Tubiphytes*-Limestone  
4  
5 1943 Member contains 51 species that belong to 39 genera, making bivalves the most diverse benthic  
6  
7 1944 macroinvertebrate group reported from this unit. However, rarefaction analysis suggests that the  
8  
9 1945 true number of bivalve species might be even higher. This bias might be partly related to the  
10  
11 1946 difficulties in the mechanical sampling of the specimens. Seven species and two genera are newly  
12  
13 1947 described, including the earliest Triassic representative of the Megalodontoida. The reported bivalve  
14  
15 1948 fauna is clearly dominated by byssally attached bivalves, which were adapted to the hard substratum  
16  
17 1949 provided by the *Tubiphytes*-microbial buildup. The biogeographic relationships are primarily  
18  
19 1950 controlled by geographic distance. This fauna is the most diverse bivalve fauna of the Bithynian in  
20  
21 1951 terms of taxonomic richness and clearly more diverse than all Early Triassic bivalve faunas. However,  
22  
23 1952 its species richness is still distinctly lower than that of the most diverse bivalve faunas from later in  
24  
25 1953 the Anisian and the Ladinian. We suggest that the studied fauna represents the beginning of a new  
26  
27 1954 recovery phase that is at least partly associated with the resurgence of organic buildups at the  
28  
29 1955 beginning of the Middle Triassic.  
30  
31  
32  
33  
34  
35 1956  
36  
37  
38 1957  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1958 *Acknowledgements:* EG is deeply grateful to the Staff of Mahmudia Quarry SC-MIN-EST SA for  
 1959 unlimited access and support along many decades in the study of the Triassic limestones and  
 1960 collecting fossils, through the persons of Director-General Eng. Lorin Poiană, Eng. Geol. Adrian Tertiş  
 1961 and Eng. Valentin Pavel. EG is greatly indebted to Dr Mircea Vişan, Dr Livia Maria Popa and Dan  
 1962 Ungureanu, who accompanied or assisted during field work in the Mahmudia Quarry for many years.  
 1963 EF is much obliged to Alexander Pohle, MSc and Lara Locatelli, BSc for numerous interesting and  
 1964 helpful discussions. This article benefited from careful reviews provided by Simon Schneider,  
 1965 Winfried Werner, Sally Thomas and one anonymous reviewer. Financial support by the Swiss  
 1966 National Science Foundation (grant 200021-162402/1 to MH) is gratefully acknowledged. Field work  
 1967 performed by EG was funded by UEFISCDI PNCDI-II-ID-1960/2009-2010.

1968

1969

#### 1970 DATA ARCHIVING STATEMENT

1971

1972 This published work and the nomenclatural acts it contains, have been registered in ZooBank:

1973 <http://zoobank.org/References/C2FC303B-A96E-4982-B9BC-977280FC42B6>

1974

1975

#### 1976 REFERENCES

1977

1978 ABERHAN, M. and MUSTER, H. 1997. Palaeobiology of Lower Jurassic Bakevelliid Bivalves from

1979 Western Canada. *Palaeontology*, **40**, 799–815.



- 1980 ALLASINAZ, A. 1966. Il Trias in Lombardia (studi geologici e paleontologici). XVIII La fauna a
- 1981 Lamellibranchi dello Julico (Carnico medio). *Rivista Italiana di Paleontologia e Stratigrafia*, **72**,
- 1982 609–752.
- 1983 — 1972. Revisione dei pettinidi Triassici. *Rivista Italiana di Paleontologia e Stratigrafia*, **78**, 189–
- 1984 428.
- 1985 ANDO, H. 1987. Paleobiological study of the Late Triassic bivalve *Monotis* from Japan. *The University*
- 1986 *Museum, The University of Tokyo, Bulletin*, **30**, 1–110.
- 1987 ASSMANN, P. 1937. Revision der Fauna der Wirbellosen der oberschlesischen Trias. *Abhandlungen*
- 1988 *der Preußischen Geologischen Landesanstalt, Neue Folge*, **170**, 134 pp., pl. 1–22.
- 1989 BALINI, M., LUCAS, S. G., JENKS, J. F. and SPIELMANN, J. A. 2010. Triassic ammonoid biostratigraphy:
- 1990 an overview. *Geological Society, London, Special Publications*, **334**, 221–262.
- 1991 BANKS, C. J. and ROBINSON, A. G. 1997. Mesozoic strike-slip back-arc basins of the Western Black Sea
- 1992 Region. In ROBINSON, A. G. (ed.) *Regional and Petroleum Geology of the Black Sea and*
- 1993 *Surrounding Region*, AAPG Memoir, 53–61.
- 1994 BAUD and RICHOZ, S. 2004. The Lower Triassic microbialites and precipitates: environmental control,
- 1995 distribution in space and time. In CAMOIN, G. and GAUTRET, P. (eds.) *Microbialites and*
- 1996 *Microbial Communities in Sedimentary Systems*, Publications AST, Paris, 15–17.
- 1997 — — and PRUSS, S. B. 2007. The lower Triassic anachronistic carbonate facies in space and time.
- 1998 *Global and Planetary Change*, **55**, 81–89.
- 1999 BEESLEY, P. L., ROSS, G. J. B. and WELLS, A. 1998. *Mollusca: The Southern Synthesis. Fauna of*
- 2000 *Australia, vol. 5, part A*. CSIRO Publishing, Melbourne, 563 pp.

- 1
- 2
- 3 2001 BEGG, J. G. and CAMPBELL, H. J. 1985. *Etalia*, a new Middle Triassic (Anisian) bivalve from New
- 4
- 5 2002 Zealand, and its relationship with other pteriomorphs. *New Zealand Journal of Geology and*
- 6
- 7 2003 *Geophysics*, **28**, 725–741.
- 8
- 9
- 10 2004 BEURLIN, K. 1944. Beiträge zur Stammesgeschichte der Muscheln.
- 11
- 12 2005 *Mathematisch-Naturwissenschaftliche Abteilung der Bayerischen Akademie der*
- 13
- 14 2006 *Wissenschaften zu München, Sitzungsberichte*, **1–2**, 133–145.
- 15
- 16
- 17
- 18 2007 — 1954. As faunas de lamellibranquios do sistema Gonduanico no Parana. *In* LANGE, W. (ed.)
- 19
- 20 2008 *Paleontologia Do Parana*, Comissão de Comemorações do Centenario do Parana, Curitiba,
- 21
- 22 2009 107–136, pl. 1–7.
- 23
- 24
- 25 2010 BIELER, R., MIKKELSEN, P. M., COLLINS, T. M., GLOVER, E. A., GONZÁLEZ, V. L., GRAF, D. L., HARPER, E.
- 26
- 27 M., HEALY, J., KAWAUCHI, G. Y., SHARMA, P. P., STAUBACH, S., STRONG, E. E., TAYLOR, J. D.,
- 28 2011
- 29 TĚMKIN, I., ZARDUS, J. D., CLARK, S., GUZMÁN, A., MCINTYRE, E., SHARP, P. and GIRIBET, G.
- 30 2012
- 31 2014. Investigating the Bivalve Tree of Life – an exemplar-based approach combining
- 32
- 33 2014 molecular and novel morphological characters. *Invertebrate Systematics*, **28**, 32–115.
- 34
- 35
- 36
- 37 2015 BITTNER, A. 1891. Triaspetrefakten von Balia in Kleinasien. *Jahrbuch der k. k. Geologischen*
- 38
- 39 *Reichsanstalt, Wien*, **41**, 97–116.
- 40 2016
- 41
- 42
- 43 2017 — 1895. Lamellibranchiaten der alpinen Trias. 1. Theil. *Abhandlungen der k. k. geologischen*
- 44
- 45 2018 *Reichsanstalt*, **18**, 1–236.
- 46
- 47
- 48 2019 — 1900. Ueber *Pseudomonotis Telleri* und verwandte Arten der unteren Trias. *Jahrbuch der k. k.*
- 49
- 50 *Geologischen Reichsanstalt*, **50**, 59–66.
- 51 2020
- 52
- 53 2021 — 1901. Lamellibranchiaten aus der Trias des Bakonyer Waldes. *Resultate der wissenschaftlichen*
- 54
- 55 *Erforschung des Balatonsees, Anhang: Palaeontologie der Umgebung des Balatonsees II*, **3**, 1–
- 56 2022
- 57 107.
- 58 2023
- 59
- 60

- 1
- 2
- 3 2024 — 1902. Brachiopoden und Lamellibranchiaten aus der Trias von Bosnien, Dalmatien und Venetien.
- 4
- 5 2025 *Jahrbuch der k. k. Geologischen Reichsanstalt*, **52** (3–4), 495–646.
- 6
- 7
- 8 2026 BLASCHKE, F. 1905. Die Gastropodenfauna der Pachycardientuffe der Seiseralpe in Südtirol nebst
- 9
- 10 2027 einem Nachtrag zur Gastropodenfauna der roten Raibler Schichten auf dem Schlernplateau.
- 11
- 12 2028 *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients*, **17**, 161–221.
- 13
- 14
- 15
- 16 2029 BÖHM, J. 1907. Über *Cardium Neptuni* Goldfuss. *Monatsberichte der Deutschen Geologischen*
- 17
- 18 2030 *Gesellschaft*, **59**, 148 pp.
- 19
- 20
- 21 2031 BOLTEN, J. F. 1798. [Published by Röding]. *Museum Boltenianum sive Catalogus Cimeliorum e Tribus*
- 22
- 23 2032 *Regnis Naturae quae olim Collegerat Joa. Fried. Bolten. Pars Secunda. Continens Conchylia sive*
- 24
- 25 2033 *Testacea Univalvia, Bivalvia, & Multivalvia*. Johan Christi Trappli, Hamburg, 199 pp.
- 26
- 27
- 28
- 29 2034 BOYD, D. W. and NEWELL, N. D. 1997. A reappraisal of Trigoniacea Families (Bivalvia) and a
- 30
- 31 2035 Description of Two New Early Triassic Species. *American Museum Novitates*, **3216**, 1–14.
- 32
- 33
- 34 2036 BRACK, P. and RIEBER, H. 1993. Toward a better definition of the Anisian/Ladinian boundary: New
- 35
- 36 2037 biostratigraphic data and correlations of boundary sections from the Southern Alps. *Eclogae*
- 37
- 38 2038 *Geologicae Helvetiae*, **86**, 415–527.
- 39
- 40
- 41
- 42 2039 — MUNDIL, R., OBERLI, F., MEIER, M. and RIEBER, H. 1996. Biostratigraphic and radiometric age
- 43
- 44 2040 data question the Milankovitch characteristics of the Latemar cycles (Southern Alps, Italy).
- 45
- 46 2041 *Geology*, **24**, 371–375.
- 47
- 48
- 49 2042 BRAYARD, A., VENNIN, E., OLIVIER, N., BYLUND, K. G., JENKS, J., STEPHEN, D. A., BUCHER, H.,
- 50
- 51 2043 HOFMANN, R., GOUEMAND, N. and ESCARGUEL, G. 2011. Transient metazoan reefs in the
- 52
- 53 2044 aftermath of the end-Permian mass extinction. *Nature Geoscience*, **4**, 693–697.
- 54
- 55
- 56
- 57 2045 — KRUMENACKER, L. J., BOTTING, J. P., JENKS, J. F., BYLUND, K. G., FARA, E., VENNIN, E., OLIVIER,
- 58
- 59 2046 N., GOUEMAND, N., SAUCÈDE, T., CHARBONNIER, S., ROMANO, C., DOGUZHAEVA, L., THUY,
- 60

- 2047 B., HAUTMANN, M., STEPHEN, D. A., THOMAZO, C. and ESCARGUEL, G. 2017. Unexpected Early  
2048 Triassic marine ecosystem and the rise of the Modern evolutionary fauna. *Science Advances*, **3**,  
2049 e1602159.
- 2050 BROGLIO LORIGA, C., NERI, C. and POSENATO, R. 1986. The Early Macrofaunas of the Werfen  
2051 Formation and the Permian-Triassic Boundary in the Dolomites (Southern Alps, Italy). *Acta*  
2052 *Geologica*, **62**, 3–18.
- 2053 — GÓCZÁN, F., HAAS, J., LENNER, K., NERI, C., ORAVECZ-SCHEFFER, A., POSENATO, R., SZABÓ, I. and  
2054 TÓTH MAKK, Á. 1990. The Lower Triassic sequences of the Dolomites (Italy) and transdanubian  
2055 mid-mountains (Hungary) and their correlation. *Memorie di Scienze Geologiche*, **42**, 41–103.
- 2056 BROILI, F. 1903. Die Fauna der Pachycardientuffe der Seiser Alp (Mit Ausschluss der Gastropoden und  
2057 Cephalopoden). *Palaeontographica*, **50**, 145–227.
- 2058 BROM, K. R., NIEDŹWIEDZKI, R., BRACHANIEC, T., FERRÉ, B. and SALAMON, M. A. 2016.  
2059 Environmental control on shell size of Middle Triassic bivalve *Plagiostoma*. *Carnets de*  
2060 *Géologie*, **16**, 297–305.
- 2061 BRONN, H. G. 1848–1849. *Index Palaeontologicus oder Übersicht der bis jetzt bekannten fossilen*  
2062 *Organismen*, bearbeitet unter Mitwirkung der Herren Prof. H. R. Göppert und Herm. v. Meyer,  
2063 bearbeitet von Dr. H. G. Bronn. Erste Abtheilung A. Nomenclator Palaeontologicus, in  
2064 Alphabetischer Ordnung. Schweizerbart. Stuttgart. 1381 pp. [1848.] Zweite Abtheilung B.  
2065 Enumerator Palaeontologicus: Systematische Zusammenstellung und Geologische  
2066 Entwicklungs-Gesetze der Organischen Reiche. 980 pp. [1849].
- 2067 BUCHER, H. 1992. Ammonoids of the Hyatti Zone and the Anisian transgression in the Triassic Star  
2068 Peak Group, north-western Nevada, USA. *Palaeontographica*, **A223**, 137–166.
- 2069 BUDAI, T. 2008. Geological excursions in the Alpine and Germanic Triassic facies areas of Hungary.  
2070 *Triassic Workshop 2008, Hungary*, 50 pp.

- 2071 BURGESS, S. D., BOWRING, S. A. and SHEN, S.-Z. 2014. High-precision timeline for Earth's most severe  
2072 extinction. *PNAS*, **111**, 3316–3321.
- 2073 BUSSE, E. 1972. Fazies und Fauna des oberen Muschelkalkes von Willebadessen. *Philippia*, **1**, 110–  
2074 126.
- 2075 CARTER, J. G. 1990. *Skeletal Biomineralisation: Patterns, Processes and Evolutionary Trends*. Volume  
2076 I, New York (Van Nostrand Rheinhold), 832 pp.
- 2077 — and HAUTMANN, M. 2011. Shell microstructure of the basal Pectinid *Pleuronectites laevigatus*:  
2078 Implications for Pectinoid phylogeny (Mollusca: Bivalvia: Pteriomorphia). *Journal of*  
2079 *Paleontology*, **85**, 464–467.
- 2080 — ALTABA, C. R., ANDERSON, L. C., ARAUJO, R., BIAKOV, A. S., BOGAN, A. E., CAMPBELL, D. D.,  
2081 CAMPBELL, M., CHEN, J.-H., COPE, J. C. W., DELVENE, G., DIJKSTRA, H. H., FANG, Z.-J.,  
2082 GARDNER, R. N., GAVRILOVA, V. A., GONCHAROVA, I. A., HARRIES, P. J., HARTMAN, J. H.,  
2083 HAUTMANN, M., HOEH, W. R., HYLLEBERG, J., BAO-YU, J., JOHNSTON, P., KIRKENDALE, L.,  
2084 KLEEMANN, K., KOPPKA, J., KŘÍŽ, J., MACHADO, D., MALCHUS, N., MÁRQUEZ-ALIAGA, A.,  
2085 MASSE, J.-P., MCROBERTS, C. A., MIDDELFART, P. U., MITCHELL, S., NEVESSKAJA, L. A., ÖZER, S.,  
2086 POJETA Jr, J., POLUBOTKO, I. V., PONS, J. M., POPOV, S., SÁNCHEZ, T., SARTORI, A. F., SCOTT, R.  
2087 W., SEY, I. I., SIGNORELLI, J. H., SILANTIEV, V. V., SKELTON, P. W., STEUBER, T., WATERHOUSE, J.  
2088 B., WINGARD, G. L. and YANCEY, T. 2011. A synoptical classification of the Bivalvia.  
2089 *Paleontological Contributions*, **4**, 1–47.
- 2090 CHEN, C. 1976. In GU, Z., HUANG, B., CHEN, C., WEN, S., MA, Q., LAN, X., XU, J., LIU, L., WANG, S.,  
2091 WANG, D., QIU, R., HUANG, Z., ZHANG, Z., CHEN, J. and WE, P. (eds.) *The Lamellibranch Fossils*  
2092 *of China*. Nanjing Institute of Geology and Palaeontology, Academia Sinica. Science Press,  
2093 Beijing, 522 pp., 150 pl.

- 2094 COHEN, K. M., FINNEY, S. C., GIBBARD, P. L. and FAN, J.-X. 2013. The ICS International
- 2095 Chronostratigraphic Chart. *Episodes*, **36**, 199–204.
- 2096 CONRAD, T. A. 1862. Descriptions of new genera, subgenera, and species of Tertiary and recent
- 2097 shells. *Proceedings of the Academy of Natural Science of Philadelphia*, **14**, 284–291.
- 2098 CONTI, M. A. and MONARI, S. 1992. Thin-shelled bivalves from the Jurassic Rosso Ammonitico and
- 2099 Calcari a *Posidonia* Formations of the Umbrian-Marchean Apennine (Central Italy).
- 2100 *Paleopelagos*, **2**, 193–213.
- 2101 COOPER, M. R. 1991. Lower Cretaceous Trigonioidea (Mollusca, Bivalvia) from the Algoa basin, with a
- 2102 revised classification of the order. *Annals of the South African Museum*, **100** (1), 1–52.
- 2103 COSSMANN, M. 1897. Paléoconchologie et ouvrages généraux. *Revue Critique de Paléozoologie*, **1**,
- 2104 46–68.
- 2105 — 1904. Note sur l'Infralias de La Vendée et des Deux- Sèvres. II. Pélécypodes. *Bulletin de la Société*
- 2106 *Géologique de France (série 4)*, **3**, 497–537.
- 2107 COX, L. R. 1929. A synopsis of the Lamellibranchia and Gastropoda of Portland beds of England, Part
- 2108 1. Lamellibranchia. *Proceedings of the Dorset Natural History and Archaeology Society*, **50**,
- 2109 131–202.
- 2110 — 1954. Taxonomic notes on Isognomonidae and Bakevelliidae. *Proceedings of the Malacological*
- 2111 *Society of London*, **31**, 46–49.
- 2112 — 1962. New genera and subgenera of Mesozoic bivalvia. *Palaeontology*, **4**, 592–598.
- 2113 — 1964. Notes concerning the taxonomy and nomenclature of fossil Bivalvia (mainly Mesozoic).
- 2114 *Proceedings of the Malacological Society of London*, **36**, 39–48.
- 2115 — NEWELL, N. D., BRANSON, C. C., CASEY, R., CHAVAN, A., COOGAN, A. H., DECHASEAUX, C.,
- 2116 FLEMING, C. A., HAAS, F., HERTLEIN, L. G., KEEN, A. M., LAROCQUE, A., MCALESTER, A. L.,

- 2117 PERKINS, B. F., PURI, H. S., SMITH, L. A., SOOT-RYEN, T., STENZEL, H. B., TURNER, R. D. and
- 2118 WEIR, J. 1969. Systematic Descriptions. In MOORE, R. C. (ed.) *Treatise on Invertebrate*
- 2119 *Paleontology, Part N, Mollusca 6, Bivalvia*, The Geological Society of America & the University
- 2120 of Kansas, Boulder, Colorado & Lawrence, Kansas, 1–952 pp.
- 2121 DALL, W. H. 1889. On the hinge of pelecypods and its development, with an attempt toward a better
- 2122 subdivision of the group. *American Journal of Science and Arts (series 3)*, **38**, 445–462.
- 2123 — 1898. *Contributions to the Tertiary fauna of Florida with especial reference to the Miocene Silex*
- 2124 *beds of Tampa and the Pliocene beds of the Caloosahatchie River, including in many cases a*
- 2125 *complete revision of the generic groups treated and of their American Tertiary Species. Part IV.*
- 2126 *1. Prionodesmacea: Nucula to Julia. 2. Teleodesmacea: Teredo to Ervilia*. Wagner Free Institute
- 2127 of Science, Philadelphia, Transactions 3(4), i-viii + p. 571–947, pl. 23–35.
- 2128 DIENER, C. 1923. Lamellibranchiata triadica. In DIENER, C. (ed.) *Fossilium Catalogus I: Animalia*, W.
- 2129 Junk, Berlin, 257 pp.
- 2130 DILLWYN, L. W. 1817. A descriptive datalogue of recent shells, arranged according to the Linnean
- 2131 method; with particular attention to the synonymy. John & Arthur Arch London, Vol. 1, 1–580,
- 2132 Vol. 2, 581–1092 + 29 pp.
- 2133 DOBRE-POPA, L. M. 2011. Evolution of microfacial domain in the eastern part (Mahmudia-Murighiol-
- 2134 Dunavătu Zone) of the Triassic carbonate platform in the Tulcea Unit, North Dobrogean
- 2135 Orogen. PhD thesis, University of Bucharest, 220 pp. [in Romanian]
- 2136 DOEDERLEIN, L. 1829. *Lateinische Synonyme und Etymologien, Dritter Theil*. Leipzig, 346 pp.
- 2137 DUFF, K. L. 1975. Palaeoecology of a bituminous shale – The Lower Oxford Clay of Central England.
- 2138 *Palaeontology*, **18**, 443–482.

- 2139 DUN, W. S. 1932. The lower marine forms of *Myonia* with notes on the proposed new genus  
2140 *Pachymyonia*. *Records of the Australian Museum*, **18**, 411–414.
- 2141 DUNKER, W. 1849. Über die im Kasseler Muschelkalk bis jetzt gefundenen Mollusken. *Programm der*  
2142 *höheren Gewerbeschule in Cassel*, 11–28.
- 2143 — 1851. Über die im Muschelkalk von Oberschlesien bis jetzt gefundenen Mollusken.  
2144 *Palaeontographica*, **1**, 283–310.
- 2145 ETTER, W. 1996. Pseudoplanktonic and benthic invertebrates in the Middle Jurassic Opalinum Clay,  
2146 northern Switzerland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **126**, 325–341.
- 2147 FÉRUSSAC, A. E. J. d'Audebard de. 1821–1822. *Tableaux Systématiques des Animaux Mollusques*  
2148 *suivis d'un Prodrome Général pour tous les Mollusques Terrestres ou Fluviatiles Vivants ou*  
2149 *Fossiles. Première Partie, Tableaux Systématiques Généraux*. Arthus-Bertrand, Paris, J.B.  
2150 Sowerby, 110 pp.
- 2151 FISCHER, P. 1880–1887. *Manuel de Conchyliologie et de Paléontologie Conchyliologique, ou Histoire*  
2152 *Naturelle des Mollusques Vivants et Fossiles suivi d'un Appendice sur les Brachiopodes par D. P.*  
2153 *Oehlert. Avec 23 planches contenant 600 figures dessinées par S. P. Woodward*. Savy, Paris.  
2154 xxiv + 1369 pp., 23 pl., 1138 text fig. (1:1–112 [1880]; 2:113–192 [1881]; 3:193–304 [1881];  
2155 4:305–416 [1882]; 5:417–512 [1883]; 6:513–608 [1883]; 7:609–688 [1884a]; 8:689–784 [29,  
2156 1885]; 9:785–896 [1885]; 10:897–1008 [1886]; 11:1009–1369 [1887]).
- 2157 FLEMING, J. 1828. *A History of British Animals, Exhibiting the Descriptive Characters and Systematical*  
2158 *Arrangement of the Genera and Species of Quadrupeds, Birds, Reptiles, Fishes, Mollusca, and*  
2159 *Radiata of the United Kingdom; including the Indigenous, Extirpated, and Extinct Kinds,*  
2160 *Together with Periodical and Occasional Visitants*. Bell & Bradfute, Edinburgh, & James  
2161 Duncan, London, 565 pp.



1  
2  
3 2162 FLÜGEL, E. 1994. Pangean shelf carbonates: Controls and paleoclimatic significance of Permian and  
4  
5 2163 Triassic reefs. *Geological Society of America Special Papers*, **288**, 247–266.  
6  
7  
8 2164 — 2002. Triassic reef patterns. *Special Publication - Society for Sedimentary Geology*, **72**, 391–463.  
9  
10  
11 2165 — and STANLEY, G. D. Jr 1984. Reorganisation, development and evolution of post-Permian reefs  
12  
13 and reef organisms. *Paleontographica Americana*, **54**, 177–186.  
14 2166  
15  
16 2167 FOREL, M.-B. and GRĂDINARU, E. 2018. First report of ostracods (Crustacea) associated with  
17  
18 Bithynian (Anisian, Middle Triassic) Tubiphytes-microbial reef in the North Dobrogean Orogen  
19 2168  
20 (Romania). *Papers in Palaeontology*, **4**, 211–244.  
21 2169  
22  
23  
24 2170 FOSTER, W. J. and SEBE, K. 2017. Recovery and diversification of marine communities following the  
25  
26 late Permian mass extinction event in the western Palaeotethys. *Global and Planetary Change*,  
27 2171  
28 **155**, 165–177.  
29 2172  
30  
31  
32 2173 — DANISE, S., PRICE, G. D. and TWITCHETT, R. J. 2017. Subsequent biotic crises delayed marine  
33  
34 recovery following the late Permian mass extinction event in northern Italy. *PLoS ONE*, **12**,  
35 2174  
36 e0172321.  
37 2175  
38  
39 2176 — — SEDLACEK, A., PRICE, G. D., HIPS, K. and TWITCHETT, R. J. 2015. Environmental controls on  
40  
41 the post-Permian recovery of benthic, tropical marine ecosystems in western Palaeotethys  
42 2177  
43 (Aggtelek Karst, Hungary). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **440**, 374–394.  
44 2178  
45  
46  
47 2179 — LEHRMANN, D. J., HIRTZ, J. A., WHITE, M., YU, M., LI, J. and MARTINDALE, R. C. 2019. Early  
48  
49 Triassic benthic invertebrates from the Great Bank of Guizhou, South China: Systematic  
50 2180  
51 palaeontology and palaeobiology. *Papers in Palaeontology*, published online 21 March 2019.  
52 2181  
53 doi: 10.1002/spp2.1252.  
54 2182  
55  
56  
57  
58  
59  
60

- 2183 FRECH, F. 1904. Neue Zweischaler und Brachiopoden aus der Bakonyer Trias. *Resultate Der*
- 2184 *Wissenschaftlichen Erforschung Des Balatonsees, Anhang: Palaeontologie Der Umgebung des*
- 2185 *Balatonsees I*, **1** (1), 1–137.
- 2186 — 1905. Nachträge zu den Cephalopoden und Zweischalern der Bakonyer Trias (Werfener und
- 2187 Cassianer Estherienschichten). *Resultate der Wissenschaftlichen Erforschung des Balatonsees*,
- 2188 **1** (1), 1–30.
- 2189 — 1909. Die Leitfossilien der Werfener Schichten und Nachträge zur Fauna des Muschelkalkes der
- 2190 Cassianer und Raibler Schichten sowie des Rhaet und des Dachsteindolomites (Hauptdolomit).
- 2191 *Resultate der Wissenschaftlichen Erforschung des Balatonsees, Anhang: Palaeontologie der*
- 2192 *Umgebung des Balatonsees*, **1** (1), 1–95.
- 2193 FRIESENBICHLER, E., RICHOSZ, S., BAUD, A., KRISTYN, L., SAHAKYAN, L., VARDANYAN, S., PECKMANN,
- 2194 J., REITNER, J. and HEINDEL, K. 2018. Sponge-microbial build-ups from the lowermost Triassic
- 2195 Chanakhchi section in southern Armenia: Microfacies and stable carbon isotopes.
- 2196 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **490**, 653–672.
- 2197 — HAUTMANN, M., NÜTZEL, A., URLICHS, M. and BUCHER, H. 2019. Palaeoecology of Late Ladinian
- 2198 (Middle Triassic) benthic faunas from the Schlern/Sciliar and Seiser Alm/Alpe di Siusi area
- 2199 (South Tyrol, Italy). *Paläontologische Zeitschrift*, **93**, 1–29.
- 2200 FÜRSICH, F. T. and WENDT, J. 1977. Biostratigraphy and palaeoecology of the cassian formation
- 2201 (Triassic) of the Southern Alps. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **22**, 257–
- 2202 323.
- 2203 GARCIA-MARCH, J. R., MÁRQUEZ-ALIAGA, A. and CARTER, J. G. 2008. The duplivincular ligament of
- 2204 recent *Pinna nobilis* L., 1758: further evidence for pterineid ancestry of the Pinnoidea. *Journal*
- 2205 *of Paleontology*, **82**, 621–627.

- 2206 GEINITZ, H. B. 1842. Über einige Petrefakten des Zechsteins und Muschelkalks. *Neues Jahrbuch für*  
 2207 *Mineralogie*, 576–579.
- 2208 GIRIBET, G. and DISTEL, D. L. 2003. Bivalve phylogeny and molecular data. *In* Lydeard, C. and  
 2209 Lindberg, D. R. (eds). *Molecular Systematics and Phylogeography of Mollusks, no 3*.  
 2210 Smithsonian Books, Washington D. C. & London, 45–90.
- 2211 GOLDFUSS, A. 1833–1841. *Petrefacta Germaniae tam ea, quae in Museo Universitatis Regiae*  
 2212 *Borussicae Fidericiae Wilhelmae Rhenanae Servantur quam alia Quaecunquae in Museis*  
 2213 *Hoenighusiano Muensteriano Aliisque Extant, Iconibus et Descriptionibus Illustrata*.  
 2214 *Abbildungen und Beschreibungen der Petrefacten Deutschlands und der Angrenzender Länder*  
 2215 *unter Mitwirkung des Herrn Grafen Georg zu Münster*. Arnz & Co. Düsseldorf. Zweiter Theil  
 2216 [Bivalvia], part 1, pp. 1–68, with Atlas, pl. 72–96 [1833]; part 2, pp. 69–140, with Atlas. pl. 97–  
 2217 121 [1835]; part 3, pp. 141–224, with Atlas, pl. 122–146 [1837]; part 4, pp. 225–312, with  
 2218 Atlas, pl. 147–165 [1841].
- 2219 GÖTZ, A. E., SZULC, J. and FEIST-BURKHARDT, S. 2005. Distribution of sedimentary organic matter in  
 2220 Anisian carbonate series of S Poland: evidence of third-order sea-level fluctuations.  
 2221 *International Journal of Earth Sciences (Geologische Rundschau)*, **94**, 267–274.
- 2222 GRĂDINARU, E. 1988. Jurassic sedimentary rocks and bimodal volcanics of the Cîrjelari-Camena  
 2223 Outcrop Belt: evidence for a transtensile regime of the Peceneaga-Camena Fault. *Studii și*  
 2224 *Cercetări de Geologie, Geofizică, Geografie, Seria Geologie*, **33**, 97–121.
- 2225 — 1995. Mesozoic rocks in North Dobrogea: an overview. *In* SĂNDULESCU, M. and GRĂDINARU, E.  
 2226 (eds.) *IGCP Project No. 369, Comparative Evolution of Peri Tethyan Rift Basin, Central and*  
 2227 *North Dobrogea, Romania, October 1–4, 1995. Field Guidebook*, Bucharest, 1–4.
- 2228 — 2000. Introduction to the Triassic geology of North Dobrogea Orogen - An overview of the  
 2229 Triassic System in the Tulcea Unit and the ammonoid biostratigraphy. *In* GRĂDINARU, E. (ed.)

- 2230 *Workshop on the Lower-Middle Triassic (Olenekian-Anisian) Boundary, 7–10 June 2000, Tulcea,*  
 2231 *Romania, Conference and Field Trip. Field Trip Guide, Bucharest, 1–63.*
- 2232 — and GAETANI, M. 2019. Upper Spathian to Bithynian (Lower to Middle Triassic) brachiopods from  
 2233 North Dobrogea (Romania). *Rivista Italiana di Paleontologia e Stratigrafia*, **125** (1), 91–123.
- 2234 GRANT-MACKIE, J. A. 1980. Mode of life and adaptive evolution in the cosmopolitan Triassic bivalve  
 2235 *Monotis*. *Journal of the Malacological Society of Australia*, **4**, 242–243.
- 2236 GRAY, J. E. 1842. Molluscs. In [multiauthored] *Eastern Zoological Gallery*. In [multiauthored], *Synopsis*  
 2237 *of the Contents of the British Museum, 44<sup>nd</sup> edn*, Woodfall and Son. London, 48–92.
- 2238 — 1847. A list of the genera of Recent Mollusca, their synonyma and types. *Zoological Society of*  
 2239 *London, Proceedings*, **15**, 129–219.
- 2240 — 1854. A revision of the arrangement of the families of bivalve shells (Conchifera). *Annals and*  
 2241 *Magazine of Natural History (series 2)*, **13** (77), 408–418; **14** (79), 21–28.
- 2242 GREGORIO, A. de 1886. Monographie des fossiles de Valpore (Mont Grappa) du sous-horizon Grappin  
 2243 de Greg. *Annales de Géologie et de Paléontologie*, **2**, 1–20.
- 2244 GU, Z., HUANG, B., CHEN, C., WEN, S., MA, Q., LAN, X., XU, J., LIU, L., WANG, S., WANG, D., QUI, R.,  
 2245 HUANG, Z., ZHANG, Z., CHEN, J. and WU, P. 1976. *The Lamellibranch Fossils of China*. Nanjing  
 2246 Institute of Geology and Palaeontology, Academia Sinica, Science Press, Beijing, 522 pp., 150  
 2247 pl.
- 2248 GUO, F. 1985. *Fossil bivalves of Yunnan*. Yunnan Science and Technology Publishing House, Kunming,  
 2249 319 pp. [in Chinese, English summary]
- 2250 HAGDORN, H. 2015. Wirbellose des Lettenkeupers. In HAGDORN, H., SCHOCH, R. and SCHWEIGERT,  
 2251 G. (eds.) *Der Lettenkeuper - Ein Fenster in die Zeit vor den Dinosauriern -*  
 2252 *Palaeodiversity-Sonderband*, Stuttgart & Ingelfingen (Schmidt), 107–140.

- 2253 — and MUNDLOS, R. 1982. Autochthonschille im Oberen Muschelkalk (Mitteltrias)
- 2254 Südwestdeutschlands. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, **162**,
- 2255 332–351.
- 2256 — and OCKERT, W. 1993. *Encrinus liliiformis* im Trochitenkalk Süddeutschlands. In HAGDORN, H.
- 2257 and SEILACHER, A. (eds.) *Muschelkalk. Sonderbände Der Gesellschaft Für Naturkunde in*
- 2258 *Württemberg e.V. 2*, Goldschneck-Verlag Werner K. Weidert, Stuttgart, Kolb, 245–260.
- 2259 — and NITSCH, E. 2009. The Germanic Triassic in its Southwest German type area. *6th International*
- 2260 *Triassic Field Workshop (Pan-European Correlation of the Triassic) Triassic of Southwest*
- 2261 *Germany 175th Anniversary of the Foundation of the Triassic System by Friedrich von Alberti*
- 2262 *September 7–11, 2009, Tübingen and Ingelfingen*, 72 pp.
- 2263 HALLAM, A. 1981. The end-Triassic bivalve extinction event. *Palaeogeography, Palaeoclimatology,*
- 2264 *Palaeoecology*, **35**, 1–44.
- 2265 HAMMER, Ø., HARPER, D. A. T. and RYAN, P. D. 2001. PAST: Paleontological Statistics Software
- 2266 Package for Education and Data Analysis. *Palaeontologia Electronica*, **4**, 1–9.
- 2267 HAUSMANN, I. M. and NÜTZEL, A. 2015. Diversity and palaeoecology of a highly diverse Late Triassic
- 2268 marine biota from the Cassian Formation of north Italy. *Lethaia*, **48**, 235–255.
- 2269 HAUTMANN, M. 2001a. Die Muschelfauna der Nayband-Formation (Obertrias, Nor-Rhät) des
- 2270 östlichen Zentraliran. *Beringeria*, **29**, 181 pp.
- 2271 — 2001b. Taxonomy and phylogeny of cementing Triassic bivalves (families Prospondylidae,
- 2272 Plicatulidae, Dimyidae and Ostreidae). *Palaeontology*, **44**, 339–373.
- 2273 — 2003. Notes on *Gruenewaldia* WÖHRMANN, 1889 (Bivalvia, Trigonioidea, Triassic). *Neues*
- 2274 *Jahrbuch für Geologie und Paläontologie Monatshefte*, **3**, 145–152.

- 2275 — 2008. Taxonomy and phylogeny of the Triassic bivalve families Mysidiellidae Cox, 1964 and  
2276 Healeyidae new family. *Journal of Paleontology*, **82**, 555–564.
- 2277 — 2010. The first scallop. *Paläontologische Zeitschrift*, **84**, 317–322.
- 2278 — and HAGDORN, H. 2013. Oysters and oyster-like bivalves from the Middle Triassic Muschelkalk of  
2279 the Germanic Basin. *Paläontologische Zeitschrift*, **87**, 19–32.
- 2280 — SMITH, A. B., MCGOWAN, A. J. and BUCHER, H. 2013. Bivalves from the Olenekian (Early Triassic)  
2281 of south-western Utah: systematics and evolutionary significance. *Journal of Systematic*  
2282 *Palaeontology*, **11** (3), 263–293.
- 2283 — BAGHERPOUR, B., BROSE, M., FRISK, Å. M., HOFMANN, R., BAUD, A., NÜTZEL, A., GOUEMAND,  
2284 N. and BUCHER, H. 2015. Competition in slow motion: The unusual case of benthic marine  
2285 communities in the wake of the end-Permian mass extinction. *Palaeontology*, **58**, 871–901.
- 2286 HAYAMI, I. 1969. Notes on Mesozoic ‘planktonic’ bivalves. *Journal of the Geological Society of Japan*,  
2287 **75**, 375–385.
- 2288 — 1975. *A systematic survey of the Mesozoic Bivalvia from Japan*. The University Museum, The  
2289 University of Tokyo, Bulletin, **10**, 249 pp.
- 2290 HODGES, P. 2000. The Early Jurassic Bivalvia from the Hettangian and Lower Sinemurian of south-  
2291 west Britain; Part 1. *Monograph of the Palaeontological Society of London, United Kingdom*,  
2292 **154**, 1–64.
- 2293 HOFMANN, R., HAUTMANN, M. and BUCHER, H. 2013a. A new paleoecological look at the Dinwoody  
2294 Formation (Lower Triassic, western USA): intrinsic versus extrinsic controls on ecosystem  
2295 recovery after the end-Permian mass extinction. *Journal of Paleontology*, **87**, 854–880.

- 2296 — — WASMER, M. and BUCHER, H. 2013*b*. Palaeoecology of the Spathian Virgin Formation (Utah,  
 2297 USA) and its implications for the Early Triassic recovery. *Acta Palaeontologica Polonica*, **58**,  
 2298 149–173.
- 2299 — — BRAYARD, A., NÜTZEL, A., BYLUND, K. G., JENKS, J. F., VENNIN, E., OLIVIER, N. and BUCHER,  
 2300 H. 2014. Recovery of benthic marine communities from the end-Permian mass extinction at  
 2301 the low latitudes of eastern Panthalassa. *Palaeontology*, **57**, 547–589.
- 2302 — — and BUCHER, H. 2015. Recovery dynamics of benthic marine communities from the Lower  
 2303 Triassic Werfen Formation, northern Italy. *Lethaia*, **48**, 474–496.
- 2304 HYATT, A. 1892. Remarks on the Pinnidae. *Proceedings, Boston Society of Natural History*, **25**, 335–  
 2305 346.
- 2306 JATTIOT, R., BRAYARD, A., BUCHER, H., VENNIN, E., CARAVACA, G., JENKS, J. F., BYLUND, K. G. and  
 2307 ESCARGUEL, G. 2018. Palaeobiogeographical distribution of Smithian (Early Triassic) ammonoid  
 2308 faunas within the Western USA Basin and its controlling parameters. *Palaeontology*, published  
 2309 online 7 June 2018. doi.org/10.1111/pala.12375.
- 2310 JEFFERIES, R. P. S. and MINTON, P. 1965. The mode of life of two Jurassic species of ‘*Posidonia*’  
 2311 (*Bivalvia*). *Palaeontology*, **8**, 156–185.
- 2312 KAIM, A. and SCHNEIDER, S. 2012. A conch with a collar: Early ontogeny of the enigmatic fossil  
 2313 bivalve *Myoconcha*. *Journal of Paleontology*, **86**, 652–658.
- 2314 KING, W. 1844. On a new genus of Palaeozoic shells. *Annals and Magazine of Natural History*, **14**,  
 2315 313–317.
- 2316 — 1850. *A Monograph of the Permian Fossils of England*. Palaeontographical Society Monograph,  
 2317 London, 258 pp., 28 pl.

- 2318 KIPARISOVA, L. D. 1938. The Lower Triassic pelecypods of the Ussuriland. *Trudy Geologicheskogo*  
 2319 *Instituta*, **7**, 197–311. [in Russian]
- 2320 KITTL, E. 1903. Geologie der Umgebung von Sarajevo. *Jahrbuch der k. k. Geologischen Reichsanstalt*,  
 2321 **53**, liii + pp. 515–748, xxi–xxiii, 1 map.
- 2322 — 1908. Beiträge zur Kenntnis der Triasbildungen der nordöstlichen Dobrudscha. *Denkschriften der*  
 2323 *Kaiserlichen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, **81**,  
 2324 447–532.
- 2325 KLIPSTEIN, A. VON 1843–1845. *Beiträge zur geologischen Kenntniss der östlichen Alpen*. Georg  
 2326 Friedrich Heyer's Verlag, Giessen, 312 pp., 20 pl.
- 2327 KLOTZ, W. and LUKAS, V. 1988. Bioherme im Unteren Muschelkalk (Trias) Südhessens. *Neues*  
 2328 *Jahrbuch für Geologie und Paläontologie Monatshefte*, **11**, 661–669.
- 2329 KOMATSU, T., HUYEN, D. T. and HUU, N. D. 2010. Radiation of Middle Triassic bivalve: Bivalve  
 2330 assemblages characterized by infaunal and semi-infaunal burrowers in a storm- and wave-  
 2331 dominated shelf, An Chau Basin, North Vietnam. *Palaeogeography, Palaeoclimatology*,  
 2332 *Palaeoecology*, **291**, 190–204.
- 2333 — CHEN, J.-H., CAO, M. Z., STILLER, F. and NARUSE, H. 2004. Middle Triassic (Anisian) diversified  
 2334 bivalves: Depositional environments and bivalve assemblages in the Leidapo Member of the  
 2335 Qingyan Formation, southern China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **208**,  
 2336 207–223.
- 2337 KOPPKA, J. 2018. Pinnidae (Bivalvia) from the Reuchenette Formation (Kimmeridgian, Upper Jurassic)  
 2338 of northwestern Switzerland. *Acta Geologica Polonica*, **68**, 21–51.
- 2339 KUTASSY, A. 1934. *Pachydonta mesozoica. Fossilium Catalogus I*. Gravenhage, **68**, 1–202.



- 2340 LAMARCK, J. B. P. A. de Monet de. 1799. Prodrome d'une nouvelle classification des coquilles,  
 2341 comprenant une rédaction appropriée des caractères génériques, et l'établissement d'un  
 2342 grand nombre de genres nouveaux. *Société d'Histoire Naturelle de Paris, Mémoires*, **1**, 63–91.
- 1809. *Philosophie Zoologique, ou Exposition des Considérations Relative à l'Histoire Naturelle des Animaux; à la Diversité de Leur Organisation et des Facultés qu'ils en Obtiennent; aux Causes Physiques qui Maintiennent en Eux la Vie et Donnent lieu aux Mouvements qu'ils Exécutent; enfin, à celles qui Produisent, les Unes le Sentiment, et les Autres l'Intelligence de Ceux qui en sont Doués*. Chez D. (ed.), L'Auteur, Paris, 2 Vols: Vol. 1, xxv + 422 pp., + Table des Matières, pp. 423–428; Vol. 2, 466 pp., + Table des Matières, pp. 467–475.
- 2349 LAUBE, G. C. 1865. *Die Fauna der Schichten von St. Cassian: Ein Beitrag zur Paläontologie der alpinen Trias, II. Abtheilung, Brachiopoden und Bivalven*. Denkschriften der Mathematisch-Naturwissenschaftlichen Classe der kaiserlich-königlichen Akademie der Wissenschaften, 25, 76 pp. + pl. 11–20.
- 2353 LEACH, W. E. 1819. Descriptions des nouvelles espèces d'animaux découvertes par le vaisseau Isabelle dans un voyage au pôle boréal. *Journal de Physique, de Chimie, d'Histoire Naturelle et des Arts*, **88**, 462–467.
- 2356 LI, M., HUANG, C., HINNOV, L., CHEN, W., OGG, J. and TIAN, W. 2018. Astrochronology of the Anisian stage (Middle Triassic) at the Guandao reference section, South China. *Earth and Planetary Science Letters*, **482**, 591–606.
- 2359 LIEBERMAN, H. M. 1979. Die Bivalven- und Ostracodenfauna von Raibl und ihr stratigraphischer Wert. *Verhandlungen der Geologischen Bundesanstalt*, **1979** (2), 85–131.
- 2361 LINNAEUS, C. A. 1758. *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. Editio Decima, reformata. Tomus I, Regnum Animale*. 823 pp.

- 2364 MANFRIN, S., MIETTO, P. and PRETO, N. 2005. Ammonoid biostratigraphy of the Middle Triassic  
2365 Latemar platform (Dolomites, Italy) and its correlation with Nevada and Canada. *Geobios*, **38**,  
2366 477–504.
- 2367 MARTINDALE, R. C., FOSTER, W. J. and VELLEDEITS, F. 2019. The survival, recovery, and diversification  
2368 of metazoan reef ecosystems following the end-Permian mass extinction event.  
2369 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **513**, 100–115.
- 2370 MARWICK, J. 1953. Divisions and Faunas of the Hokonui System. *New Zealand Geological Survey*,  
2371 *Palaeontological Bulletin*, **21**, 141 pp.
- 2372 MCGOWAN, A. J., SMITH, A. B. and TAYLOR, P. D. 2009. Faunal diversity, heterogeneity and body size  
2373 in the Early Triassic: Testing post-extinction paradigms in the Virgin Limestone of Utah, USA.  
2374 *Australian Journal of Earth Sciences*, **56**, 859–872.
- 2375 MCROBERTS, C. A. 2010. Biochronology of Triassic bivalves. In LUCAS, S. G. (ed.) *The Triassic*  
2376 *Timescale*, Geological Society, London, Special Publication, 334, 201–219.
- 2377 — 2017. A Silicified Late Triassic (Norian) Bivalve Molluscan Fauna from the Alexander Terrane,  
2378 Southeastern Alaska. *Bulletins of American Paleontology*, **391**, 1–110.
- 2379 — KRYSTYN, L. and SHEA, A. 2008. Rhaetian (Late Triassic) *Monotis* (Bivalvia: Pectinoida) from the  
2380 Eastern Northern Calcareous Alps (Austria) and the End-Norian crisis in pelagic faunas.  
2381 *Palaeontology*, **51**, 721–735.
- 2382 MEEK, F. B. 1865. Description of fossils from the auriferous slates of California. *Geological Survey of*  
2383 *California, Geology*, **1**, 477–482.
- 2384 — and HAYDEN, F. V. 1865. Palaeontology of the Upper Missouri. *Smithsonian Contributions to*  
2385 *Knowledge*, **14**, 1–135.

1  
2  
3 2386 — and WORTHEN, A. H. 1866. Descriptions of Paleozoic fossils from the Silurian, Devonian, and  
4  
5 2387 Carboniferous rocks of Illinois and other western states. *Chicago Academy of Natural Science,*  
6  
7 2388 *Proceedings*, **1**, 11–23.  
8  
9  
10 2389 MILLER, S. A. 1877. *The American Palaeozoic Fossils, a Catalog of the Genera and Species, with names*  
11  
12 *of authors, dates, places of publication, groups of rock in which found, and the etymology and*  
13 2390  
14 *signification of the words, and an introduction devoted to the stratigraphical geology of the*  
15 2391  
16 *Palaeozoic rocks*. Published by the author, Cincinnati, 253 pp.  
17 2392  
18  
19  
20 2393 MIRĂUȚĂ, E. and IORDAN, M. 1982. New Triassic fossiliferous localities in the Tulcea Zone (North  
21  
22 Dobrogea). *Dări de Seamă, Institutul de geologie și geofizică*, **66**, 15–22.  
23 2394  
24  
25  
26 2395 — — and GHEORGHIAN, D. 1984. New biostratigraphic data on the Triassic from the Somova-  
27  
28 2396 Sarica Hill area (Tulcea Zone, North Dobrogea). *Dări de Seamă, Institutul de geologie și*  
29  
30 2397 *geofizică*, **68**, 35–48.  
31  
32  
33 2398 MOJSISOVICS, E. 1874. Über die Triadischen Pelecypoden-Gattungen *Daonella* und *Halobia*.  
34  
35 2399 *Abhandlungen der k.k. Geologischen Reichsanstalt*, **7**, 1–38.  
36  
37  
38 2400 MONARI, S. 2003. A new genus and species of fimbriid bivalve from the Kimmeridgian of the Western  
39  
40 Pontides, Turkey, and the phylogeny of the Jurassic Fimbriidae. *Palaeontology*, **46**, 857–884.  
41 2401  
42  
43  
44 2402 MONNET, C. and BUCHER, H. 2006. Anisian (Middle Triassic) ammonoids from North America:  
45  
46 2403 quantitative biochronology and biodiversity. *Stratigraphy*, **2**, 281–296.  
47  
48  
49 2404 MORRIS, J. and LYCETT, J. 1853. *Monograph of the Mollusca from the Great Oolite, chiefly from*  
50  
51 2405 *Minchinhampton and the coast of Yorkshire*. Palaeontographical Society (Monographs),  
52  
53 London. Part II, Bivalves: 1–80, pl. 1–8. Part III, Bivalves: 81–147, pl. 9–15.  
54 2406  
55  
56 2407 MURCHISON, R. I., BUCKMAN, J. and STRICKLAND, H. A. 1845. Outline of the Geology of the  
57  
58 Neighborhood of Cheltenham. *John Murray*, London, 109 pp., 13 pl.  
59 2408  
60

- 2409 MUSTER, H. 1995. Taxonomie und Paläobiogeographie der Bakevelliidae (Bivalvia). *Beringeria*, **14**,  
 2410 161 pp.
- 2411 MÜLLER, O. F. 1776. *Zoologiae Danicae Prodomus, seu Animalium Daniae et Norvegiae Indigenarum*  
 2412 *Characteres, Nomina, et Synonyma Imprimis Popularium*. Impensis Auctoris. Hallageriis. Havniae  
 2413 (Copenhagen), 282 pp.
- 2414 MÜNSTER, G. 1841. Beiträge zur Geognosie und Petrefacten-Kunde des südöstlichen Tirols vorzüglich  
 2415 der Schichten von St. Cassian. II. Beschreibung und Abbildung der in den Kalkmergelschichten  
 2416 von St. Cassian gefunden Versteinerungen. *Beiträge zur Petrefacten-Kunde*, **4**, 25–152.
- 2417 NAWROCKI, J. and SZULC, J. 2000. The Middle Triassic magnetostratigraphy from the Peri-Tethys  
 2418 basin in Poland. *Earth and Planetary Science Letters*, **182**, 77–92.
- 2419 NERI, C. and POSENATO, R. 1985. New biostratigraphical data on uppermost Werfen Formation of  
 2420 western Dolomites (Trento, Italy). *Geologische und Paläontologische Mitteilungen Innsbruck*,  
 2421 **14**, 83–107.
- 2422 NEUMAYR, M. 1884. Zur Morphologie des Bivalvenschlosses. *Sitzungsberichte der Kaiserlichen*  
 2423 *Akademie der Wissenschaften in Wien*, **88**, 385–418.
- 2424 — 1891. Beiträge zu einer morphologischen Eintheilung der Bivalven. *Denkschriften der Kaiserlichen*  
 2425 *Akademie der Wissenschaften, Wien, Mathematisch-Naturwissenschaftliche Classe*, **58**, 701–  
 2426 801.
- 2427 NEWELL, N. D. 1965. Classification of the Bivalvia. *American Museum Novitates*, **2206**, 1–25.
- 2428 — and BOYD, D. W. 1975. Parallel Evolution in Early Trigoniacean Bivalves. *Bulletin of the American*  
 2429 *Museum of Natural History*, **154**, 53–162.
- 2430 — — 1987. Iteration of ligament structures in Pteriomorphian Bivalves. *American Museum*  
 2431 *Novitates*, **2875**, 1–111.

- 2432 NEYMAN, A. A. 1967. Limits to the application of the trophic group concept in benthic studies.
- 2433 *Oceanology*, **7**, 149–155.
- 2434 NÜTZEL, A. and KAIM, A. 2014. Diversity, palaeoecology and systematics of a marine fossil
- 2435 assemblage from the Late Triassic Cassian Formation at Settsass Scharte, N Italy.
- 2436 *Paläontologische Zeitschrift*, **88**, 405–431.
- 2437 — — and GRĂDINARU, E. 2018. Middle Triassic (Anisian, Bithynian) gastropods from North
- 2438 Dobrogea (Romania) and their significance for gastropod recovery from the end-Permian mass
- 2439 extinction event. *Papers in Palaeontology*, published online 6 May 2018.
- 2440 doi:10.1002/spp2.1115.
- 2441 OCKERT, W. 1993. Die Zwergfaunaschichten (Unterer Hauptmuschelkalk, Trochitenkalk, mo1) im
- 2442 nordöstlichen Baden-Württemberg. In HAGDORN, H. and SEILACHER, A. (eds.) *Muschelkalk.*
- 2443 *Sonderbände der Gesellschaft für Naturkunde in Württemberg e.V.* 2, 117–215.
- 2444 OGILVIE GORDON, M. M. 1927. Das Grödener-, Fassa- und Enneberggebiet in den Südtiroler
- 2445 Dolomiten. *Abhandlungen der Geologischen Bundesanstalt*, **24**, I und II Teil, 376 pp., III Teil, 89
- 2446 pp.
- 2447 OKAY, A. I., ŞENGÖR, A. M. C. and GÖRÜR, N. 1994. Kinematic history of the opening of the Black Sea
- 2448 and its effect on the surrounding regions. *Geology*, **22**, 267–270.
- 2449 ORBIGNY, A. d' 1849. *Prodrome de Paléontologie stratigraphique universelle des animaux mollusques*
- 2450 *et rayonnés*. Faisant Suite au Cours Élémentaire de Paléontologie et de Geologie
- 2451 Stratigraphiques, vol. 1, Victor Masson, Paris, 394pp.
- 2452 PARONA, C. F. 1889. Studio monografico della fauna raibliana di Lombardia. *Memoria Premiata dal*
- 2453 *Regio Istituto Lombardo di Scienze e Lettere*, 156 pp.

- 2454 PETERS, K. F. 1867. Grundlinien zur Geographie und Geologie der Dobrudscha. *Denkschriften der*  
2455 *kaiserlichen Akademie der Wissenschaften*, **27**, 83–207.
- 2456 PHILLIPS, J. 1829. *Illustrations of the Geology of Yorkshire; or a Description of the Strata and Organic*  
2457 *Remains of the Yorkshire Coast: Accompanied by a Geological Map, Sections, and Plates of the*  
2458 *Fossil Plants and Animals*. Thomas Wilson & Sons, York, 192 pp. + 14 pl.
- 2459 PIA, J. 1937. *Stratigraphie und Tektonik der Pragser Dolomiten in Südtirol*. Eigenverlag, Wien.
- 2460 PICARD, E. 1907. Die Gattung *Pinna* in der Trias. *Jahrbuch der Königlich Preussischen Geologischen*  
2461 *Landesanstalt und Bergakademia zu Berlin für das Jahr 1903*, **24**, 333–336.
- 2462 POPA, L., PANAIOTU, C. E. and GRĂDINARU, E. 2014. An early Middle Anisian (Middle Triassic)  
2463 *Tubiphytes* and cement crusts-dominated reef from North Dobrogea (Romania): Facies,  
2464 depositional environment and diagenesis. *Acta Geologica Polonica*, **64**, 189–206.
- 2465 POSENATO, R. 2008. Anisian (Middle Triassic) bivalves from the Dolomites (Italy). *Neues Jahrbuch für*  
2466 *Geologie und Paläontologie Abhandlungen*, **247** (1), 93–115.
- 2467 QIHONG, M., CHEN, J., LAN, X., GU, Z., CHEN, C. and LIN, M. 1976. Mesozoic lamellibranchs from  
2468 Yunnan. In *Nanjing Institute of Geology and Palaeontology, Academia Sinica (ed.), Mesozoic*  
2469 *Fossils from Yunnan, Vol. 1*, Science Press, Beijing, 161–386 pp. [in Chinese]
- 2470 QUENSTEDT, F. A. 1851–1852. *Handbuch der Petrefaktenkunde*. Laupp and Siebeck. Tübingen.  
2471 Volume 1, 1–528, Volume 2, 529–792.
- 2472 RAFINESQUE, C. S. 1815. *Analyse de la Nature, ou Tableau de l'Univers et des Corps Organisés, etc.* C.  
2473 S. Rafinesque. Jean Barravecchia. Palermo, 223 pp.
- 2474 RAUP, D. M. 1979. Size of the Permo-Triassic bottleneck and its evolutionary implications. *Science*,  
2475 217–218.

- 2476 REIS, O. M. 1926. Die Fauna des Wettersteinkalks. III Theil: Gastropoden, Bivalven, Brachiopoden.  
2477 *Geognostisches Jahreshft (München), Jahrgang 39*, 87–188.
- 2478 RIDING, R. and GUO, L. 1992. Affinities of *Tubiphytes*. *Palaeontology*, **35**, 37–49.
- 2479 ROS-FRANCH, S., MÁRQUEZ-ALIAGA, A. and DAMBORENEA, S. 2014. Comprehensive database on  
2480 Induan (Early Triassic) to Sinemurian (Early Jurassic) marine bivalve genera and their  
2481 paleobiogeographic record. *University of Kansas Paleontological Contributions*, **8**, 1–219.
- 2482 ROSSI RONCHETTI, C. 1959. Il Trias in Lombardia: studi geologici e paleontologici: Lamellibranchi  
2483 ladinici del Gruppo delle Grigne. *Istituto di geologia, paleontologia e geografia fisica della*  
2484 *Università di Milano*, **65**, 269–357.
- 2485 RUNNEGAR, B. 1974. Evolutionary history of the bivalve subclass Anomalodesmata. *Journal of*  
2486 *Paleontology*, **48**, 904–939.
- 2487 SACCHI VIALLI, G. and VAI, A. 1958. Revisione della fauna triassica bresciana. La fauna dell'Anisico.  
2488 *Atti dell'Istituto Geologica della Università di Pavia*, **8**, 41–91.
- 2489 SALOMON, W. 1895. Geologische und palaeontologische Studien über die Marmolata.  
2490 *Palaeontographica*, **42**, 1–210.
- 2491 SANDERS, H. F. 1968. Marine benthic diversity: a comparative study. *The American Naturalist*, **102**,  
2492 243–282.
- 2493 SĂNDULESCU, M. 1980. Analyse géotectonique des chaines alpines situées autour de la Mer Noire  
2494 occidentale. *Anuarul Institutului de Geologie și Geofizică*, **56**, 5–54.
- 2495 — 1995. Dobrogea within the Carpathian Foreland. In SĂNDULESCU, M. and GRĂDINARU, E. (eds.)  
2496 *IGCP Project No. 369, Comparative Evolution of Peri Tethyan Rift Basin, Central and North*  
2497 *Dobrogea, Romania, October 1–4, 1995. Field Guidebook*, Bucharest, 1–4.

- 2498 SCHAUROTH, K. VON 1859. Kritisches Verzeichniss der Versteinerungen der Trias im Vicentinischen.
- 2499 *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien,*
- 2500 *Mathematisch-Naturwissenschaftliche Klasse*, **34**, 283–356.
- 2501 SCHLAGER, W. 2000. Sedimentation rates and growth potential of tropical, cool-water and mud-
- 2502 mound carbonate factories. In INSALACO, E., SKELTON, P. W. and PALMER, T. J. (eds.)
- 2503 *Carbonate Platform Systems: Components and Interactions*, Geological Society, London, Special
- 2504 Publication, **178**, London, 217–227.
- 2505 — 2003. Benthic carbonate factories of the Phanerozoic. *International Journal of Earth Sciences*, **92**,
- 2506 445–464.
- 2507 SCHLOTHEIM, E. F. VON 1820. *Die Petrefaktenkunde auf ihrem jetzigen Standpunkte durch die*
- 2508 *Beschreibung seiner Sammlung versteinerner und fossiler Überreste des Thier- und*
- 2509 *Pflanzenreichs der Vorwelt erläutert*. Becker, Gotha, 437 pp.
- 2510 — 1823. *Nachträge zur Petrefaktenkunde*. Zweite Abtheilung. Becker, Gotha, 114 pp.
- 2511 SCHMIDT, M. 1928. *Die Lebewelt unserer Trias*. Hohenlohesche Buchhandlung F. Rau, Öhringen.
- 2512 — 1938. *Die Lebewelt unserer Trias - Nachtrag*. Hohenlohesche Buchhandlung F. Rau, Öhringen.
- 2513 SCHNEIDER, S. 2017. *Pachyrisma grande* Morris & Lycett, 1850: redescription and assessment of its
- 2514 potential role as an ancestor to the Hippuritida. *Zitteliana*, **89**, 271–284.
- 2515 SCOPOLI, J. A. 1777. *Introductio ad historiam naturalem sistens genera Lapidum, Plantarum, et*
- 2516 *Animalium hactenus detecta, characteribus essentialibus donata, in Tribus divisa, subinde ad*
- 2517 *leges naturae*. Wolfgangum Gerle, Prague, 506 + 36 pl.
- 2518 SEILACHER, A. 1954. Ökologie der Triassischen Muschel *Lima lineata* (Schloth.) und ihrer Epöken.
- 2519 *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*, **1954** (4), 163–183.



- 2520 ŞENGÖR, A. M. C. 1984. The Cimmeride Orogenic System and the tectonics of Eurasia. *Geological*
- 2521 *Society of America Special Paper*, **195**, 82 pp.
- 2522 — 1986. Die Alpiden und die Kimmeriden: die verdoppelte Geschichte der Tethys. *Geologische*
- 2523 *Rundschau*, **75**, 501–510.
- 2524 SENOWBARI-DARYAN, B., ZÜHLKE, R., BECHSTÄDT, T. and FLÜGEL, E. 1993. Anisian (Middle Triassic)
- 2525 Buildups of the Northern Dolomites (Italy): The Recovery of Reef Communities after the
- 2526 Permian/Triassic Crisis. *Facies*, **28**, 181–256.
- 2527 SHEVYREV, A. A. 1995. *Triassic ammonites of northwestern Caucasus*. Nauka, Moscow, 179 pp. [in
- 2528 Russian]
- 2529 SHIGETA, Y., ZAKHAROV, Y. D., MAEDA, H. and POPOV, A. M. 2009. The Lower Triassic System in the
- 2530 Abrek Bay area, South Primorye, Russia. *National Museum of Nature and Science Monographs*,
- 2531 **38**, 1–218.
- 2532 SILBERLING, N. J. and NICHOLS, K. M. 1982. Middle Triassic molluscan fossils of biostratigraphic
- 2533 significance from the Humboldt Range, northwestern Nevada. *Geological Survey Professional*
- 2534 *Paper*, **1207**, 1–77.
- 2535 — GRANT-MACKIE, J. A. and NICHOLS, K. M. 1997. The Late Triassic bivalve *Monotis* in accreted
- 2536 terranes of Alaska. *U. S. Geological Survey Bulletin*, **2115**, 1–21.
- 2537 SIMIONESCU, I. 1908. Über das Vorkommen der Werfener Schichten in Dobrogea (Rumänien).
- 2538 *Verhandlungen der k. k. geologischen Reichsanstalt*, **7**, 159–161.
- 2539 — 1910a. Studii geologice şi paleontologice din Dobrogea. III. Fauna triasică dela Deşli-Căira. La
- 2540 faune triasique de Deşli-Căira (Dobrogea). *Academia Română, Publicaţiunile Fondului Vasile*
- 2541 *Adamachi*, **5**, 495–524.

- 2542 — 1910b. Studii geologice și paleontologice din Dobrogea. IV. Fauna triasică din Insula Popina.
- 2543 *Academia Română, Publicațiunile Fondului Vasile Adamachi*, **5**, 495–524.
- 2544 — 1911. Studii geologice și paleontologice din Dobrogea. V. Fauna triasică inferioară din Dobrogea
- 2545 (La faune du Trias inférieur de Dobrogea). *Academia Română, Publicațiunile Fondului Vasile*
- 2546 *Adamachi*, **5**, 63–79.
- 2547 — 1925. Păturile cu Daonella in Dobrogea (Les couches à Daonella de Dobrogea). *Academia*
- 2548 *Română, Publicațiunile Fondului Vasile Adamachi*, **9**, 1–10.
- 2549 SIMPSON, E. H. 1949. Measurement of diversity. *Nature*, **163**, 688.
- 2550 SOWERBY, J. 1814. *The Mineral Conchology of Great Britain; or Coloured Figures and Description of*
- 2551 *those Remains of Testaceous Animals or Shells Which Have Been Preserved at Various Times*
- 2552 *and Depths in the Earth*. Benjamin Meredith. London, **1** (9–14), 97–178, pl. 45–78.
- 2553 — 1819. *The Mineral Conchology of Great Britain; or Coloured Figures and Description of those*
- 2554 *Remains of Testaceous Animals or Shells Which Have Been Preserved at Various Times and*
- 2555 *Depths in the Earth*. Benjamin Meredith, London, **3** (39–44), 41–98, pl. 222–253.
- 2556 — 1824. *The Mineral Conchology of Great Britain; or Coloured Figures and Description of those*
- 2557 *Remains of Testaceous Animals or Shells Which Have Been Preserved at Various Times and*
- 2558 *Depths in the Earth*. Benjamin Meredith, London, **5** (77–83), 65–138, pl. 444–485.
- 2559 SPECIALE, A. 1967. Fossili del Trias Medio delle Valli Trompia e Sabbia. *Rivista Italiana di*
- 2560 *Paleontologia e Stratigrafia*, **73**, 1055–1140.
- 2561 STANLEY, G. D. Jr 1988. The History of Early Mesozoic reef communities: A three step process.
- 2562 *Palaios*, **3**, 170–183.
- 2563 STANLEY, S. M. 1970. Relation of shell form to life habits of the Bivalvia (Mollusca). *Geological Society*
- 2564 *of America, Memoir*, **125**, 1–296.

- 2565 — 1972. Functional Morphology and Evolution of Byssally Attached Bivalve Mollusks. *Journal of*  
2566 *Paleontology*, **46**, 165–212.
- 2567 — 2016. Estimates of the magnitudes of major marine mass extinctions in earth history.  
2568 *Proceedings of the National Academy of Sciences*, **113**, E6325–E6334.
- 2569 STAROBOGATOV, Y. 1992. Morphological basis for phylogeny and classification of Bivalvia. *Ruthenica*,  
2570 **2**, 1–25.
- 2571 STILLER, F. 2001. Fossilvergesellschaftungen, Paläökologie und paläosynökologische Entwicklung im  
2572 Oberen Anis (Mittlere Trias) von Qingyan, insbesondere Bangtoupou, Provinz Guizhou,  
2573 Südwestchina. *Münstersche Forschung zur Geologie und Paläontologie*, **92**, 1–523.
- 2574 — and CHEN, J. 2006. New Mysidiellidae (Bivalvia) from the Anisian (Middle Triassic) of Qingyan,  
2575 South-West China. *Palaeontology*, **49**, 213–227.
- 2576 — and BUCHER, H. 2008. Anisian ammonoids from Qingyan, southwestern China: Biostratigraphical  
2577 implications for the age of the Qingyan Formation. *Swiss Journal of Geosciences*, **101**, 547–562.
- 2578 STOLICZKA, F. 1870–1871. The Pelecypoda, with a review of all known genera of this class, fossil and  
2579 Recent. In OLDHAM, T. (ed.) *Cretaceous Fauna of Southern India. Vol. 3, Palaeontologia Indica*,  
2580 *Being Figures and Descriptions of the Organic Remains Procured During the Progress of the*  
2581 *Geological Survey of India*, Trübner and Co. Memoirs of the Geological Survey of India,  
2582 Calcutta, (serie 1) 3:xxii + 535 p., 50 pl. Issued in 13 parts: dates of issue: **1–4**, 1–222, pl. 1–12  
2583 [September 1, 1870]; **5–8**, 223–409, pl. 13–28 [March 1, 1871]; **9–13**, 410–535, pl. 29–50  
2584 [August 1, 1871].
- 2585 STOPPANI, A. 1858–1860. *Les petrifications d’Esino ou description des fossiles appartenants au depot*  
2586 *superieur des environs d’Esino en Lombardie*. Paléontologie Lombarde **1**, 1–151 + 32 pl.

- 2587 — 1865. Géologie et Paléontologie des Couches à *Avicula contorta* en Lombardie. Paléontologie  
 2588 Lombarde III, Milano, 1–267.
- 2589 STROMBECK, A. VON 1849. Über *Cucullaea Beyrichii* aus dem Muschelkalke. *Zeitschrift der Deutschen*  
 2590 *Geologischen Gesellschaft*, **1**, pp. 451.
- 2591 SZABÓ, I., KOVÁCS, S., LELKES, G. and ORAVECZ-SCHEFFER, A. 1979. Stratigraphic investigation of a  
 2592 (triassic) Pelsonian-Fassanian section at Felsőörs (Balaton Highland, Hungary). *Rivista Italiana*  
 2593 *di Paleontologia e Stratigrafia*, **85**, 789–806.
- 2594 SZENTE, I., VÖRÖS, A. 2003. The Pelsonian Bivalvia Fauna of the Balaton Highland. *Geologica*  
 2595 *Hungarica*, **55**, 123–137.
- 2596 TEPPNER, W. VON 1922. Lamellibranchia Tertiaria, Anisomyaria II. In DIENER, C. (ed.) *Fossilium*  
 2597 *Catalogus, I. Animalia, Pars*, W. Junk, Berlin, 67–296.
- 2598 TOKUYAMA, A. 1959. Late Triassic Pteriacea from the Atsu and Mine Series, West Japan. *Japanese*  
 2599 *journal of geology and geography*, **30**, 1–19.
- 2600 TOMMASI, A. 1911. I fossili della lumachella triasica di Ghegna in Valsecca presso Roncobello.  
 2601 *Memorie di Paleontologia*, **27**, 1–36.
- 2602 TURCULET, I. 2009. New data about the Ammonite fauna from the Triassic developed in the Wengen  
 2603 facies at Cataloi (North Dobruja Orogen). *Geologie Tomul LV*, **2**, 45–50.
- 2604 TURNER, R. D. and ROSEWATER, J. 1958. The family Pinnidae in the western Atlantic, Johnsonia.  
 2605 *Department of Mollusks. Museum of Comparative Zoology, Harvard University*, **3**, 285–326.
- 2606 URLICH, M. 1982. Zur Stratigraphie und Fossilführung des Lettenkeupers (Trias) bei Schwäbisch Hall  
 2607 (Baden-Württemberg). *Jahresberichte und Mitteilungen des Oberrheinischen Geologischen*  
 2608 *Vereins, Neue Folge*, **64**, 213–224.

- 2609 — 1992. Vermeintlicher Zwergwuchs bei Muscheln aus dem untersten Trochitenkalk (Oberer  
 2610 Muschelkalk, Mitteltrias) Württembergs. *Carolinea*, **50**, 9–26.
- 2611 — 2014. A phylogenetic lineage of *Myophoria kefersteini* subspecies (Bivalvia) from the Late  
 2612 Ladinian to the Early Carnian (Triassic) in the Dolomites and Julian Alps (Italy). *Neues Jahrbuch  
 2613 für Geologie und Paläontologie, Abhandlungen*, **274**, 187–207.
- 2614 VÉGH-NEUBRANDT, E. 1982. *Triassische Megalodontaceae Entwicklung, Stratigraphie und  
 2615 Paläontologie*. Akadémiai Kiadó, Budapest, 526 pp.
- 2616 VISARION, M., SĂNDULESCU, M., ROȘCA, V., STĂNICĂ, D. and ATANASIU, L. 1990. La Dobrogea dans  
 2617 le cadre de l'avant pays carpatique. *Revue Roumaine de Géophysique*, **34**, 55–65.
- 2618 VOKES, H. E. 1967. Genera of the Bivalvia: A systematic and bibliographic catalogue. *Bulletins of  
 2619 American Paleontology*, **51** (232), [103–111] + 112–394.
- 2620 — 1980. *Genera of the Bivalvia: A systematic and Bibliographic Catalogue (Revised and Updated)*.  
 2621 Paleontological Research Institution, Ithaca, 307pp.
- 2622 VU KHUC, 1991. *Paleontological Atlas of Vietnam. Volume 3. Mollusca*. Science and Technics  
 2623 Publishing House, Hanoi, 207 pp., 67 pl.
- 2624 — 2000. The Triassic of Indochina peninsula and its interregional correlation. In YIN, H., DICKINS, J.  
 2625 M., SHI, G. R. and TONG, J. (eds). *Permian-Triassic Evolution of Tethys and Western Circum-  
 2626 Pacific*. Elsevier, 392 pp.
- 2627 — and HUYEN, D. T. 1984. Some new Bivalve genera and subgenera from Triassic sediments of  
 2628 North Vietnam. *Journal Cac khoa hoc ve Trai dat*, **6** (4), Hanoi. [in Vietnamese]
- 2629 WAAGEN, L. 1906. Über die Lamellibranchiaten der Frommbachtuffe nebst Bemerkungen über deren  
 2630 verwandtschaftliche Beziehungen. *Verhandlungen der k. k. geologischen Reichsanstalt*, **16**,  
 2631 385–395.

- 2632 — 1907. Die Lamellibranchiaten der Pachycardientuffe der Seiser Alm nebst vergleichend  
2633 paläontologischen und phylogenetischen Studien. Als Fortsetzung (II. Teil) zu A. Bittner,  
2634 Lamellibranchiaten der alpinen Trias. *Abhandlungen der k. k. geologischen Reichsanstalt*, **18**,  
2635 1–180.
- 2636 WAGNER, R. 1897. Beitrag zur genaueren Kenntnis des Muschelkalks bei Jena. *Abhandlungen der*  
2637 *Königlich Preussischen Geologischen Landesanstalt Neue Folge*, **27**, 1–105.
- 2638 WALLER, T. R. 1978. Morphology, morphoclines and a new classification of the Pteriomorpha  
2639 (Mollusca: Bivalvia). *Philosophical Transaction of the Royal Society B1*, **284**, 345–365.
- 2640 — 1984. The ctenolium of scallop shells: functional morphology and evolution of a key family-level  
2641 character in the Pectinacea (Mollusca: Bivalvia). *Malacologia*, **25**, 203–219.
- 2642 — and STANLEY, G. D. Jr 2005. Middle Triassic Pteriomorphian Bivalvia (Mollusca) from the New  
2643 Pass Range, West-Central Nevada: Systematics, Biostratigraphy, Paleoecology and  
2644 Paleobiogeography. *Journal of Paleontology*, **79** (1), 64 pp.
- 2645 WASMER, M., HAUTMANN, M., HERMANN, E., WARE, D., ROOHI, G., UR-REHMAN, K., YASEEN, A. and  
2646 BUCHER, H. 2012. Olenekian (Early Triassic) bivalves from the Salt Range and Surghar Range,  
2647 Pakistan. *Palaeontology*, **55**, 1043–1073.
- 2648 WATERHOUSE, J. B. 2001. Late Paleozoic Brachiopoda and Mollusca chiefly from Wairaki Downs, New  
2649 Zealand, with notes on Scyphozoa and Triassic ammonoids and new classifications of  
2650 Linoproductoidea (Brachiopoda) and Pectinida (Bivalvia). *Earthwise*, **3**, 1–196.
- 2651 WHITFIELD, R. P. 1885. Brachiopoda and Lamellibranchiata of the Raritan clays and greensand marls  
2652 of New Jersey. *U.S. Geological Survey Monograph*, **9**, 338 pp.
- 2653 WHITTAKER, R. H. 1965. Dominance and diversity in land plant communities. *Science*, **147**, 250–260.

- 2654 WILCKENS, R. 1909. Paläontologische Untersuchung triadischer Faunen aus der Umgebung von  
2655 Predazzo in Südtirol. *Verhandlungen des Naturhistorisch-Medizinischen Vereins Heidelberg*, **10**,  
2656 81–231.
- 2657 WINKLER, G. G. 1859. *Die Schichten der Avicula contorta inner- und ausserhalb der Alpen*. Joh. Palm's  
2658 Hofbuchhandlung, München, 50 pp.
- 2659 WISSMANN, H. L. 1841. Taxonomic names, in Beiträge zur Geognosie und Petrefacten-Kunde des  
2660 südöstlichen Tirols vorzüglich der Schichten von St. Cassian. *Beiträge zur Petrefactenkunde*, **4**,  
2661 1–152.
- 2662 WITTENBURG, P. W. VON 1908. Beiträge zur Kenntnis der Werfener Schichten Südtirols.  
2663 *Abhandlungen der Geologie und Paläontologie (new series)*, **8** (5), 44 pp.
- 2664 — 1909. Einige Lamellibranchiaten der Salt-Range, mit Berücksichtigung der Lamellibranchiaten des  
2665 Süd-Ussuri-Gebiets. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, **1909** (1), 6–  
2666 13.
- 2667 WÖHRMANN, S. VON 1889. Die Fauna der sogenannten Cardita- und Raibler-Schichten in den  
2668 nordtiroler und bayerischen Alpen. *Jahrbuch der k. k. Geologischen Reichsanstalt*, **39**, 181–258,  
2669 pl. 5–10.
- 2670 — and KOKEN, E. 1892. Die Fauna der Raibler Schichten vom Schlernplateau. *Zeitschrift der*  
2671 *Deutschen geologischen Gesellschaft*, **44**, 167–223.
- 2672 YIN, H. and YIN, J. 1983. Bivalves. In YANG, Z., YIN, H., XU, G., Wu, S., HE, Y., LIU, G. and Yin, J. (eds.)  
2673 *Triassic of the South Qilian Mountains*, Geological Publishing House, Beijing, 128–175.
- 2674 YIN, J. and MCROBERTS, C. A. 2006. Latest Triassic-Earliest Jurassic bivalves of the Germig Formation  
2675 from Lanongla (Tibet, China). *Journal of Paleontology*, **80**, 104–120.
- 2676 ZAPFE, H. 1957. Dachsteinkalk und 'Dachsteinmuscheln'. *Natur und Volk*, **87**, 87–94.

- 2677 ZARDINI, R. 1981. *Fossili Cassiani (Trias Medio-Superiore). Atlante dei Bivalvi della Formazione di S.*  
 2678 *Cassiano raccolti nella regione Dolomitica attorno a Cortina d'Ampezzo*. Edizione Ghedina,  
 2679 Cortina d'Ampezzo, 1–14 + 40 pl.

# FIGURE CAPTIONS

**FIG. 1.** A, tectonic map of the North Dobrogea Orogen. Inset map showing the regional geological setting and the location of the North Dobrogea Orogen. *Abbreviations:* CU = Consul Unit; MU = Măcin Unit; NU = Niculițel Unit; TU = Tulcea Unit. B, geological map of the Mahmudia region. C, field photograph showing the Bithynian sediments of the Caerace Formation. The dashed line marks the boundary between the Stromatactis-Limestone Member and the overlying *Tubiphytes*-Limestone Member. Car (in circle) for scale. D, close-up view showing the position of the sample site. Modified after Grădinaru (2000), Popa *et al.* (2014), Forel & Grădinaru (2018), Nützel *et al.* (2018) and Grădinaru & Gaetani (2019).

**FIG. 2.** Stratigraphic column of the Mahmudia Quarry. Numbers 99–122 represent sample levels from previous works (Nützel *et al.* 2018; Grădinaru & Gaetani 2019). The star marks the sample level that yielded the bivalve specimens described herein.

**FIG. 3.** A–B, *Parallelodon?* sp., LPB IIL 2001; A, LV; B, dorsal view of A showing the chevron grooves of the duplivincular ligament (arrow). C–D, *Modiolus* sp. A, LPB IIL 2002; C, LV; D, anterior view of C. E, LV of *Modiolus* sp. B, LPB IIL 2004. F, RV of *Bositra wengensis* (Wissmann, 1841), LPB IIL 2008. G,



1  
2  
3 2700 RV of *Pinna simionescui* n. sp., holotype, LPB IIL 2006. H, LV of *Atrina multicostata* n. sp., holotype,  
4  
5 2701 LPB IIL 2005. I, not oriented valve of *Pinna*? cf. *raibliana* Parona, 1889, LPB IIL 2007. J–K, LVs of  
6  
7 2702 *Pteria sturi* (Bittner, 1895), LPB IIL 2009 and LPB IIL 2010. L, RV of *Pteria* sp. A, LPB IIL 2013. M, RV  
8  
9 2703 of *Pteria* sp. B, LPB IIL 2014. N, RV’s interior of *Pteria* cf. *cassiana* (Bittner, 1895), LPB IIL 2012. Scale  
10  
11 2704 bars represent 1 cm (A–D, F–N) and 0.5 cm (E).  
12  
13  
14  
15 2705  
16  
17  
18 2706 **FIG. 4.** A, LV of *Cultriopsis* sp., LPB IIL 2015. B–C, *Gervillaria subelegans* (Chen, 1976), LPB IIL 2016; B,  
19  
20 2707 LV; C, umbonal region of B. D, LV of *Bakevelliidae* indet., LPB IIL 2018. E–G, *Aviculopectinoidea* n.  
21  
22 2708 gen. et n. sp.; E, RV, the deep byssal notch and the sculpture of the auricles are well visible, LPB IIL  
23  
24 2709 2021; F, LV, LPB IIL 2022; G, RV, LPB IIL 2023. H, LV of ‘*Pecten*’ *volaris* (Bittner, 1902), LPB IIL 2019. I,  
25  
26 2710 LV of *Asoella*? sp., LPB IIL 2038. J–K, LVs of *Amphijanira*? aff. *landrana* (Bittner, 1895), LPB IIL 2024  
27  
28 2711 and LPB IIL 2025. L–N, *Neomorphotis compta* (Goldfuss, 1833); L and N; RVs, LPB IIL 2028 and LPB IIL  
29  
30 2712 2029; M; LV, LPB IIL 2027. Scale bars represent 1 cm.  
31  
32  
33  
34  
35 2713  
36  
37  
38 2714 **FIG. 5.** A–B, LVs of *Leptochondria pervulgata* (Bittner, 1902), LPB IIL 2039 and LPB IIL 2040. C–D, LVs  
39  
40 2715 of *Leptochondria separata* (Reis, 1926), LPB IIL 2049 and LPB IIL 2050. E, LV of *Ornithopecten* cf.  
41  
42 2716 *katzeri* (Bittner, 1902), LPB IIL 2058. F–H LVs of *Leptochondria viezzenensis* (Wilckens, 1909), LPB IIL  
43  
44 2717 2053, LPB IIL 2054 and LPB IIL 2055. I, LV of *Ornithopecten interruptus* (Bittner, 1902), LPB IIL 2057.  
45  
46 2718 J–M, *Chlamys* (*Praechlamys*) *prima* n. sp.; J and L, LVs, the auricle’s sculpture is well visible,  
47  
48 2719 paratypes, LPB IIL 2060 and 2061; K, LV, holotype, LPB IIL 2062; M, external mould of a LV, paratype,  
49  
50 2720 LPB IIL 2063. N, RV of *Pleuronectites*? cf. *balatonicus* (Bittner, 1901), LPB IIL 2068. O, LV of  
51  
52 2721 *Avichlamys*? cf. *nicolensis* (Ogilvie Gordon, 1927), LPB IIL 2059. P–Q, Pectinidae indet.; P, internal  
53  
54 2722 mould of a RV, LPB IIL 2066; Q, internal mould of a LV, LPB IIL 2067. R, LV of *Entolium reticulatum* n.  
55  
56 2723 sp. showing the shell’s reticular sculpture, holotype, LPB IIL 2070. S–U *Entolioides deecke* (Parona,  
57  
58 2724 1889). S, LV. Arrow shows the depression subparallel to posterior margin, LPB IIL 2071. T, LV, LPB IIL

2725 2072; U, interior of a LV. Dotted line marks the posterior adductor muscle, LPB IIL 2073. V, LV of  
 2726 *Entolium* sp., LPB IIL 2159. Scale bars represent 1 cm (A–D, F, H–V) and 0.5 cm (E and G).

2727

2728 **FIG. 6.** A–D, *Scythentolium anisicum* n. sp.; A, RV of holotype, LPB IIL 2075; B, LV of holotype, LPB IIL  
 2729 2075; C, RV with *Plagiostoma striatum* (LPB IIL 2102), paratype, LPB IIL 2076; D, fragmented RV  
 2730 showing the antimarginal structure (arrow), paratype, LPB IIL 2082. E–G, RVs of *Mysidioptera*  
 2731 *forficata* Bittner, 1895, LPB IIL 2089, LPB IIL 2090 and LPB IIL 2091, the weak radial sculptural  
 2732 elements are best visible in E and G. H–J, *Plagiostoma striatum* (von Schlotheim, 1820); H, RV, LPB IIL  
 2733 2102; I, anterior view of a RV showing the lunula (arrow), LPB IIL 2103; J, RV, LPB IIL 2104. K, LV of  
 2734 *Umbrostrea?* sp., LPB IIL 2087. L, LV of an articulated specimen of *Mysidioptera* sp., LPB IIL 2101. M–  
 2735 N, *Plagiostoma* cf. *angulatum* (Münster, 1841); M, fragmented LV well showing the shell's sculpture,  
 2736 LPB IIL 2128; N, RV, LPB IIL 2127. O–P, Monotidae indet., LPB IIL 2088; O, internal mould of a LV; P,  
 2737 dorsal view of O. Scale bars represent 1 cm.

2738

2739 **FIG. 7.** A–B, *Elegantinia elegans* (Dunker, 1849), LPB IIL 2130; A, RV's exterior; B, interior of A  
 2740 showing the two merged hinge teeth (arrow). C, RV of *Heminajas?* sp., LPB IIL 2131. D, RV of  
 2741 *Joannina* aff. *joannae* Waagen, 1906, LPB IIL 2132. E, fragmented RV of *Myoconcha* sp., LPB IIL 2133.  
 2742 F–H, *Unicardium schmidii* (Geinitz, 1842); F, RV, LPB IIL 2134; G, umbonal region of H showing the  
 2743 nymph (arrow); H, LV, LPB IIL 2135. I–K, *Schafhaeutlia astartiformis* (Münster, 1841); I, RV, LPB IIL  
 2744 2137; J, dorsal view of I showing parts of the hinge (arrow); K, RV, LPB IIL 2138. L, LV of  
 2745 *Schafhaeutlia?* *trigona* (Stoppani, 1858), LPB IIL 2140. M–P, *Praedicerocardium vetulus* n. gen. et n.  
 2746 sp.; M, dorsal view of an articulated specimen, holotype, LPB IIL 2142; N, anterior view of a RV,  
 2747 paratype, LPB IIL 2143; O, dorsal view of N; P, lateral view of N. Scale bars represent 1 cm (A, C–P)  
 2748 and 0.5 cm (B).

2749

1  
2  
3 2750 **FIG. 8.** A–B, *Romaniamya mahmudiaensis* n. gen. et n. sp., holotype, LPB IIL 2145; A, LV; B, posterior  
4  
5 2751 view of A showing the sulcus located dorsally to the umbonal ridge (black arrow) and the shell's  
6  
7 2752 narrow gape (grey arrow). C, internal mould of a RV of *Anomalodesmata* indet. A, LPB IIL 2148. D,  
8  
9 2753 not oriented specimen of *Anomalodesmata* indet. B, LPB IIL 2149. E, internal mould of a LV of  
10  
11 2754 *Anomalodesmata* indet. C, LPB IIL 2147. F–H, *Mytiliconcha* cf. *orobica* Tommasi, 1911; F, RV, LPB IIL  
12  
13 2755 2155; G, umbonal region of F; H, RV, LPB IIL 2156. Scale bars represent 1 cm (A, C–H) and 0.5 cm (B).

14  
15  
16  
17 2756  
18  
19  
20 2757 **FIG. 9.** Pie chart showing the percentage of bivalve genera (innermost circle), families (second  
21  
22 2758 innermost circle), superfamilies (third innermost circle), suborders (middle circle), orders (third  
23  
24 2759 outermost circle), infraclasses (second outermost circle) and subclasses (outermost circle).  
25  
26 2760 *Abbreviations for subclasses:* i.s. = *incertae sedis*. *Abbreviations for orders:* Ar = Arcoida; Me =  
27  
28 2761 Megalodontida; Modiom = Modiomorphoidea; Trigonii = Trigoniida; ? = uncertain. *Abbreviations for*  
29  
30 2762 *suborders:* Mo = Monotidina; Os = Ostreina; ? = uncertain. *Abbreviations for superfamilies:* Ar =  
31  
32 2763 Arcoidea; K = Kalenteroidea; M = Modiomorphoidea; Me = Megalodontoidea; Mo = Monotoidea; My  
33  
34 2764 = Myophorioidea; Mytiloi = Mytiloidea; Os = Ostreoidea; Ph = Pholadomyoidea; Po = Posidonioidea; ?  
35  
36 2765 = uncertain. *Abbreviations for families:* An = Antijaniridae; Bakevelli = Bakevelliidae; Di =  
37  
38 2766 Dicerocardiidae; H = Healeyidae; He = Heteropectinidae; Ka = Kalenteridae; Lucini = Lucinidae; Ma =  
39  
40 2767 Mactromyidae; Me = Megadesmidae; Mo = Monotidae; Myo = Myophoriidae; Mytili = Mytilidae;  
41  
42 2768 Ornitho = Ornithopectinidae; Os = Ostreidae; Pa = Parallelodontidae; Pl = Pleuronectitidae; Po =  
43  
44 2769 Posidoniidae; ? = uncertain. *Abbreviations for genera:* ? = uncertain.

45  
46  
47  
48  
49  
50 2770  
51  
52  
53 2771 **FIG. 10.** Pie chart showing the percentage of species (left circle) and genera (right circle) that (i) are  
54  
55 2772 newly described, (ii) the oldest and (iii) the youngest occurrence of the taxon, (iv) presented in open  
56  
57 2773 nomenclature, (v) species uncertainly belonging to a genus that is confirmed through the presence of  
58  
59 2774 another species and (vi) species and genera that do not belong to any of the just mentioned

2775 categories. The numbers within the circular segments represent the number of species belonging to  
 2776 the corresponding category.

2777

2778 **FIG. 11.** Pie chart showing the percentage of species occurring in the Western Tethys (W Tethys), the  
 2779 Muschelkalk, the Eastern Tethys (E Tethys) and northwestern Panthalassa (NW Panthalassa) during  
 2780 the Early (E. T.) and the Middle Triassic (M. T.), as well as endemic species. The numbers within the  
 2781 circular segments represent the number of species belonging to the corresponding palaeogeographic  
 2782 province.

2783

2784 **FIG. 12.** Rank-abundance distribution and guild structure of the investigated bivalve fauna. n = total  
 2785 number of individuals.

2786

2787 **FIG. 13.** Assumed life positions of epibyssate (1–12), cementing (13), free-lying (14–19), endobyssate  
 2788 (20–24), shallow burrowing (25–28) and deep burrowing (29–32) bivalve genera with respect to the  
 2789 substratum. Abbreviations: 1 = *Plagiostoma*; 2 = *Pteria*; 3 = *Mysidioptera*; 4 = *Neomorphotis*; 5 =  
 2790 *Ornithopecten*; 6 = *Chlamys* (*Praechlamys*); 7 = '*Pecten*'; 8 = *Pleuronectites*?; 9 = *Asoella*?; 10 =  
 2791 *Avichlamys*?; 11 = *Amphijanira*?; 12 = *Aviculopectinoidea* n. gen.; 13 = *Umbrostrea*?; 14 = *Entolium*;  
 2792 15 = *Praedicerocardium* n. gen.; 16 = *Gervillaria*; 17 = *Leptochondria*; 18 = *Entolioides*; 19 =  
 2793 *Scythentolium*; 20 = *Joannina*; 21 = *Cultriopsis*; 22 = *Pinna*; 23 = *Atrina*; 24 = *Modiolus*; 25 =  
 2794 *Schafhaeutlia*; 26 = *Elegantinia*; 27 = *Heminajas*?; 28 = *Unicardium*; 29 = *Romaniamya* n. gen.; 30 =  
 2795 *Anomalodesmata* indet. A; 31 = *Anomalodesmata* indet. C; 32 = *Anomalodesmata* indet. B; 33 =  
 2796 *Tubiphytes*-microbial buildup.


2797

1  
2  
3 2798 **FIG. 14.** Bivalve species richness throughout the Early and Middle Triassic based on data shown in  
4  
5 2799 Table 1. The star marks the investigated bivalve fauna. *Abbreviations:* A = Aegean; B = Bithynian; CF =  
6  
7 2800 Cassian Formation; D = Dienerian; Fassan = Fassanian; G = Griesbachian; I = Induan; JF = Jena  
8  
9 2801 Formation; LM = Leidapo Member; Longobard = Longobardian; MF = Meissner Formation; P =  
10  
11 2802 Pachycardientuffe; Pel = Pelsonian; Sm = Smithian; TK = Trochitenkalk Formation; VF = Virgin  
12  
13 2803 Formation. Absolute ages according to Cohen *et al.* (2013; updated). Information about the relative  
14  
15 2804 duration of substages come from Nawrocki & Szulc (2000), Götz *et al.* (2005), Jattiot *et al.* (2018), Li  
16  
17 2805 *et al.* (2018). See text for further information.  
18  
19  
20  
21  
22 2806

23  
24  
25 2807 **FIG. 15.** Rarefaction analysis of the reported bivalve fauna from Dobrogea (star) compared with  
26  
27 2808 those of other Middle Triassic (M. T.) as well as Early Triassic (E. T.) bivalve faunas. Data used from  
28  
29 2809 Stiller (2001), Hofmann *et al.* (2013a, b, 2014), Foster *et al.* (2015, 2019), Hautmann *et al.* (2015),  
30  
31 2810 Foster & Sebe (2017) and Friesenbichler *et al.* (2019).  
32  
33  
34  
35 2811  
36  
37  
38 2812  
39  
40  
41


42 2813 **TABLE CAPTIONS**  
43  
44 2814  
45  
46

47 2815 **TABLE 1.** Species richness of selected Early and Middle Triassic bivalve faunas. Modified after  
48  
49 2816 Friesenbichler *et al.* (2019). Information about the age of the lithological units comes from Fürsich &  
50  
51 2817 Wendt (1977), Broglio Loriga *et al.* (1990), Brack & Rieber (1993), Brack *et al.* (1996), Szente & Vörös  
52  
53 2818 (2003), Manfrin *et al.* (2005), Stiller & Bucher (2008), Hagdorn & Nitsch (2009), Shigeta *et al.* (2009),  
54  
55 2819 Balini *et al.* (2010), Hofmann *et al.* (2013a, b, 2014, 2015), Popa *et al.* (2014), Urlichs (2014),  
56  
57 2820 Hausmann & Nützel (2015), Hautmann *et al.* (2015), Foster *et al.* (2015, 2017), Brayard *et al.* (2017),  
58  
59 2821 Foster & Sebe (2017) and Foster *et al.* (2019).  
60




**FIG. 1.** A, tectonic map of the North Dobrogea Orogen. Inset map showing the regional geological setting and the location of the North Dobrogea Orogen. *Abbreviations:* CU = Consul Unit; MU = Măcin Unit; NU = Niculițel Unit; TU = Tulcea Unit. B, geological map of the Mahmudia region. C, field photograph showing the Bithynian sediments of the Caerace Formation. The dashed line marks the boundary between the Stromatactis-Limestone Member and the overlying *Tubiphytes*-Limestone Member. Car (in circle) for scale. D, close-up view showing the position of the sample site. Modified after Grădinaru (2000), Popa *et al.* (2014), Forel & Grădinaru (2018), Nützel *et al.* (2018) and Grădinaru & Gaetani (2019).

165x171mm (300 x 300 DPI)



**FIG. 2.** Stratigraphic column of the Mahmudia Quarry. Numbers 99–122 represent sample levels from previous works (Nützel *et al.* 2018; Grădinaru & Gaetani 2019). The star marks the sample level that yielded the bivalve specimens described herein.


109x108mm (300 x 300 DPI)



**FIG. 3.** A–B, *Parallelodon?* sp., LPB IIIL 2001; A, LV; B, dorsal view of A showing the chevron grooves of the duplivincular ligament (arrow). C–D, *Modiolus* sp. A, LPB IIIL 2002; C, LV; D, anterior view of C. E, LV of *Modiolus* sp. B, LPB IIIL 2004. F, RV of *Bositra wengensis* (Wissmann, 1841), LPB IIIL 2008. G, RV of *Pinna simionescui* n. sp., holotype, LPB IIIL 2006. H, LV of *Atrina multicostata* n. sp., holotype, LPB IIIL 2005. I, not oriented valve of *Pinna?* cf. *raibliana* Parona, 1889, LPB IIIL 2007. J–K, LVs of *Pteria sturi* (Bittner, 1895), LPB IIIL 2009 and LPB IIIL 2010. L, RV of *Pteria* sp. A, LPB IIIL 2013. M, RV of *Pteria* sp. B, LPB IIIL 2014. N, RV's interior of *Pteria* cf. *cassiana* (Bittner, 1895), LPB IIIL 2012. Scale bars represent 1 cm (A–D, F–N) and 0.5 cm (E).


165x204mm (300 x 300 DPI)





**FIG. 4.** A, LV of *Cultriopsis* sp., LPB IIIIL 2015. B–C, *Gervillaria subelegans* (Chen, 1976), LPB IIIIL 2016; B, LV; C, umbonal region of B. D, LV of Bakevelliidae indet., LPB IIIIL 2018. E–G, Aviculopectinoidea n. gen. et n. sp.; E, RV, the deep byssal notch and the sculpture of the auricles are well visible, LPB IIIIL 2021; F, LV, LPB IIIIL 2022; G, RV, LPB IIIIL 2023. H, LV of ‘*Pecten*’ *volaris* (Bittner, 1902), LPB IIIIL 2019. I, LV of *Asoella?* sp., LPB IIIIL 2038. J–K, LVs of *Amphijanira?* aff. *landrana* (Bittner, 1895), LPB IIIIL 2024 and LPB IIIIL 2025. L–N, *Neomorphotis comta* (Goldfuss, 1833); L and N, RVs, LPB IIIIL 2028 and LPB IIIIL 2029; M, LV, LPB IIIIL 2027. Scale bars represent 1 cm.


165x207mm (300 x 300 DPI)



**FIG. 5.** A–B, LVs of *Leptochondria pervulgata* (Bittner, 1902), LPB IIIIL 2039 and LPB IIIIL 2040. C–D, LVs of *Leptochondria separata* (Reis, 1926), LPB IIIIL 2049 and LPB IIIIL 2050. E, LV of *Ornithopecten* cf. *katzeri* (Bittner, 1902), LPB IIIIL 2058. F–H LVs of *Leptochondria viezzenensis* (Wilckens, 1909), LPB IIIIL 2053, LPB IIIIL 2054 and LPB IIIIL 2055. I, LV of *Ornithopecten interruptus* (Bittner, 1902), LPB IIIIL 2057. J–M, *Chlamys* (*Praechlamys*) *prima* n. sp.; J and L, LVs, the auricle's sculpture is well visible, paratypes, LPB IIIIL 2060 and 2061; K, LV, holotype, LPB IIIIL 2062; M, external mould of a LV, paratype, LPB IIIIL 2063. N, RV of *Pleurnectites*? cf. *balatonicus* (Bittner, 1901), LPB IIIIL 2068. O, LV of *Avichlamys*? cf. *nicolensis* (Ogilvie Gordon, 1927), LPB IIIIL 2059. P–Q, Pectinidae indet.; P, internal mould of a RV, LPB IIIIL 2066; Q, internal mould of a LV, LPB IIIIL 2067. R, LV of *Entolium reticulatum* n. sp. showing the shell's reticular sculpture, holotype, LPB IIIIL 2070. S–U *Entolioides deeckeii* (Parona, 1889). S, LV. Arrow shows the depression subparallel to posterior margin, LPB IIIIL 2071. T, LV, LPB IIIIL 2072; U, interior of a LV. Dotted line marks the posterior adductor muscle, LPB IIIIL 2073. V, LV of *Entolium* sp., LPB IIIIL 2159. Scale bars represent 1 cm (A–D, F, H–V) and 0.5 cm (E and G).


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

165x221mm (300 x 300 DPI)




**FIG. 6.** A–D, *Scythentolium anisicum* n. sp.; A, RV of holotype, LPB IIL 2075; B, LV of holotype, LPB IIL 2075; C, RV with *Plagiostoma striatum* (LPB IIL 2102), paratype, LPB IIL 2076; D, fragmented RV showing the antimarginal structure (arrow), paratype, LPB IIL 2082. E–G, RVs of *Mysidioptera fornicata* Bittner, 1895, LPB IIL 2089, LPB IIL 2090 and LPB IIL 2091, the weak radial sculptural elements are best visible in E and G. H–J, *Plagiostoma striatum* (von Schlotheim, 1820); H, RV, LPB IIL 2102; I, anterior view of a RV showing the lunula (arrow), LPB IIL 2103; J, RV, LPB IIL 2104. K, LV of *Umbrostrea*? sp., LPB IIL 2087. L, LV of an articulated specimen of *Mysidioptera* sp., LPB IIL 2101. M–N, *Plagiostoma* cf. *angulatum* (Münster, 1841); M, fragmented LV well showing the shell's sculpture, LPB IIL 2128; N, RV, LPB IIL 2127. O–P, Monotidae indet., LPB IIL 2088; O, internal mould of a LV; P, dorsal view of O. Scale bars represent 1 cm.

165x181mm (300 x 300 DPI)




**FIG. 7.** A–B, *Elegantinia elegans* (Dunker, 1849), LPB IIL 2130; A, RV's exterior; B, interior of A showing the two merged hinge teeth (arrow). C, RV of *Heminajas?* sp., LPB IIL 2131. D, RV of *Joannina* aff. *joannae* Waagen, 1906, LPB IIL 2132. E, fragmented RV of *Myoconcha* sp., LPB IIL 2133. F–H, *Unicardium schmidii* (Geinitz, 1842); F, RV, LPB IIL 2134; G, umbonal region of H showing the nymph (arrow); H, LV, LPB IIL 2135. I–K, *Schafhaeutlia astartiformis* (Münster, 1841); I, RV, LPB IIL 2137; J, dorsal view of I showing parts of the hinge (arrow); K, RV, LPB IIL 2138. L, LV of *Schafhaeutlia? trigona* (Stoppani, 1858), LPB IIL 2140. M–P, *Praedicerocardium vetulus* n. gen. et n. sp.; M, dorsal view of an articulated specimen, holotype, LPB IIL 2142; N, anterior view of a RV, paratype, LPB IIL 2143; O, dorsal view of N; P, lateral view of N. Scale bars represent 1 cm (A, C–P) and 0.5 cm (B).

165x199mm (300 x 300 DPI)




**FIG. 8.** A–B, *Romaniamya mahmudiaensis* n. gen. et n. sp., holotype, LPB IIL 2145; A, LV; B, posterior view of A showing the sulcus located dorsally to the umbonal ridge (black arrow) and the shell's narrow gape (grey arrow). C, internal mould of a RV of *Anomalodesmata* indet. A, LPB IIL 2148. D, not oriented specimen of *Anomalodesmata* indet. B, LPB IIL 2149. E, internal mould of a LV of *Anomalodesmata* indet. C, LPB IIL 2147. F–H, *Mytiliconcha* cf. *orobica* Tommasi, 1911; F, RV, LPB IIL 2155; G, umbonal region of F; H, RV, LPB IIL 2156. Scale bars represent 1 cm (A, C–H) and 0.5 cm (B).

165x75mm (300 x 300 DPI)



## Palaeontology






**FIG. 10.** Pie chart showing the percentage of species (left circle) and genera (right circle) that (i) are newly described, (ii) the oldest and (iii) the youngest occurrence of the taxon, (iv) presented in open nomenclature, (v) species uncertainly belonging to a genus that is confirmed through the presence of another species and (vi) species and genera that do not belong to any of the just mentioned categories. The numbers within the circular segments represent the number of species belonging to the corresponding category.


165x62mm (300 x 300 DPI)






**FIG. 11.** Pie chart showing the percentage of species occurring in the Western Tethys (W Tethys), the Muschelkalk, the Eastern Tethys (E Tethys) and northwestern Panthalassa (NW Panthalassa) during the Early (E. T.) and the Middle Triassic (M. T.), as well as endemic species. The numbers within the circular segments represent the number of species belonging to the corresponding palaeogeographic province.

109x58mm (300 x 300 DPI)




**FIG. 12.** Rank-abundance distribution and guild structure of the investigated bivalve fauna.  $n$  = total number of individuals.

165x112mm (300 x 300 DPI)




**FIG. 13.** Assumed life positions of epibyssate (1–12), cementing (13), free-lying (14–19), endobyssate (20–24), shallow burrowing (25–28) and deep burrowing (29–32) bivalve genera with respect to the substratum. Abbreviations: 1 = *Plagiostoma*; 2 = *Pteria*; 3 = *Mysidioptera*; 4 = *Neomorphotis*; 5 = *Ornithopecten*; 6 = *Chlamys* (*Praechlamys*); 7 = '*Pecten*'; 8 = *Pleuronectites*?; 9 = *Asoella*?; 10 = *Avichlamys*?; 11 = *Amphijanira*?; 12 = *Aviculopectinoidea* n. gen.; 13 = *Umbrostrea*?; 14 = *Entolium*; 15 = *Praedicerocardium* n. gen.; 16 = *Gervillaria*; 17 = *Leptochondria*; 18 = *Entolioides*; 19 = *Scythentolium*; 20 = *Joannina*; 21 = *Cultriopsis*; 22 = *Pinna*; 23 = *Atrina*; 24 = *Modiolus*; 25 = *Schafhaeutlia*; 26 = *Elegantinia*; 27 = *Heminajas*?; 28 = *Unicardium*; 29 = *Romaniamya* n. gen.; 30 = *Anomalodesmata* indet. A; 31 = *Anomalodesmata* indet. C; 32 = *Anomalodesmata* indet. B; 33 = *Tubiphytes*-microbial buildup.

165x111mm (300 x 300 DPI)



**FIG. 14.** Bivalve species richness throughout the Early and Middle Triassic based on data shown in Table 1. The star marks the investigated bivalve fauna. *Abbreviations:* A = Aegean; B = Bithynian; CF = Cassian Formation; D = Dienerian; Fassin = Fassanian; G = Griesbachian; I = Induan; JF = Jena Formation; LM = Leidapo Member; Longobard = Longobardian; MF = Meissner Formation; P = Pachycardientuffe; Pel = Pelsonian; Sm = Smithian; TK = Trochitenkalk Formation; VF = Virgin Formation. Absolute ages according to Cohen *et al.* (2013; updated). Information about the relative duration of substages come from Nawrocki & Szulc (2000), Götz *et al.* (2005), Jattiot *et al.* (2018), Li *et al.* (2018). See text for further information.

165x125mm (300 x 300 DPI)



**FIG. 15.** Rarefaction analysis of the reported bivalve fauna from Dobrogea (star) compared with those of other Middle Triassic (M. T.) as well as Early Triassic (E. T.) bivalve faunas. Data used from Stiller (2001), Hofmann *et al.* (2013a, b, 2014), Foster *et al.* (2015, 2019), Hautmann *et al.* (2015), Foster & Sebe (2017) and Friesenbichler *et al.* (2018b).

165x102mm (300 x 300 DPI)

Lithological Unit	Age	Number of bivalve species	References
Cassian Fm. (Italy)	Longobardian–Julian	296	Laube (1865), Bittner (1895), Pia (1937), Fürsich & Wendt (1977), Zardini (1981), Nützel & Kaim (2014), Hausmann & Nützel (2015)
Schlernplateau beds (Italy)	late Longobardian – early Julian	27	von Wöhrmann & Koken (1892), Friesenbichler <i>et al.</i> (2019)
Pachycardientuffe (Wengen Fm., Italy)	late Longobardian	150	Broili (1903), Blaschke (1905), Waagen (1907), Fürsich & Wendt (1977), Friesenbichler <i>et al.</i> (2019)
Erfurt Fm. (‘Lower Keuper’, Germany)	early Longobardian	47	Schmidt (1928, 1938), Urlichs (1982)
Meissner Fm. (upper part of the ‘Upper Muschelkalk’, Germany)	late Illyrian – early Longobardian	118	Wagner (1897), Schmidt (1928, 1938), Busse (1972), Hagdorn & Mundlos (1982), Hagdorn & Ockert (1993)
Marmolada Limestone (Italy)	middle Illyrian – middle Longobardian	45	Salomon (1895)
Latemar Limestone (Italy)	middle Illyrian – early Longobardian	24	Wilckens (1909)
Trochitenkalk Fm. (basal part of the ‘Upper Muschelkalk’, Germany)	middle Illyrian	96	Wagner (1897), Schmidt (1928, 1938), Busse (1972), Hagdorn & Ockert (1993), Ockert (1993), Urlichs (1992), Hautmann & Hagdorn (2013)
Karlstadt Fm., Heilbronn Fm., Diemel Fm. (‘Middle Muschelkalk’, Germany)	early Illyrian – middle Illyrian	34	Schmidt (1928, 1938)
Leidapo Mb. (Quingyan Fm., China)	late Pelsonian – early Illyrian	109	Stiller (2001), Komatsu <i>et al.</i> (2004)
Zuhány Limestone (Hungary)	late Pelsonian – early Illyrian	19	Foster & Sebe (2017)
Felsőörs Fm. (Hungary)	Pelsonian – Illyrian	39	Szabó <i>et al.</i> (1979), Vörös (2003)
Jena Fm. (‘Lower Muschelkalk’, Germany)	late Bithynian – early Illyrian	115	Wagner (1897), Schmidt (1928, 1938)
Lapis Limestone (Hungary)	late Bithynian – early Pelsonian	12	Foster & Sebe (2017)
<i>Tubiphytes</i> -Limestone Mb. (Romania)	Bithynian	51	this study
Röt Fm. (‘Upper Buntsandstein’, Germany)	Early Bithynian	38	Wagner (1897), Schmidt (1928, 1938)
Viganvár Limestone (Hungary)	late Aegean – early Bithynian	10	Foster & Sebe (2017)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Csopak Marl (Hungary)	Spathian	15	Broglia Loriga <i>et al.</i> (1990), Budai (2008)
Virgin Fm. (USA)	Spathian	30	McGowan <i>et al.</i> (2009), Hautmann <i>et al.</i> (2013), Hofmann <i>et al.</i> (2013 <i>b</i> )
Szinpetri Limestone Fm. (USA)	late Spathian	6	Foster <i>et al.</i> (2015)
San Lucano Mb. (Werfen Fm., Italy)	late Spathian	7	Neri & Posenato (1985), Broglia Loriga <i>et al.</i> (1990), Foster <i>et al.</i> (2017)
Cencenighe Mb. (Werfen Fm, Italy)	middle Spathian	25	Neri & Posenato (1985), Broglia Loriga <i>et al.</i> (1990), Hofmann <i>et</i> <i>al.</i> (2015), Foster <i>et al.</i> (2017)
Szin Marl Fm. (Hungary)	early – middle Spathian	10	Foster <i>et al.</i> (2015)
Val Badia Mb. (Werfen Fm., Italy)	early Spathian	15	Broglia Loriga <i>et al.</i> (1990), Hofmann <i>et al.</i> (2015), Foster <i>et</i> <i>al.</i> (2017)
Lower Shale unit (Thaynes Gr., USA)	late Smithian – early Spathian	5	Brayard <i>et al.</i> (2017)
Sinbad Fm. (USA)	Smithian	18	Hofmann <i>et al.</i> (2014)
Hidegkút Fm. (Hungary)	Smithian	15	Bittner (1901), Broglia Loriga <i>et</i> <i>al.</i> (1990)
Campil Mb. (Werfen Fm., Italy)	Smithian	21	Broglia Loriga <i>et al.</i> (1990), Hofmann <i>et al.</i> (2015), Foster <i>et</i> <i>al.</i> (2017)
Zhitkov Fm. (Russia)	Dienerian – Smithian	11	Shigeta <i>et al.</i> (2009)
Gastropod Oolite (Werfen Fm., Italy)	late Dienerian – early Smithian	11	Broglia Loriga <i>et al.</i> (1990), Hofmann <i>et al.</i> (2015), Foster <i>et</i> <i>al.</i> (2017)
Seis Mb. (Werfen Fm., Italy)	Dienerian	18	Broglia Loriga <i>et al.</i> (1990), Hofmann <i>et al.</i> (2015), Foster <i>et</i> <i>al.</i> (2017)
Dinwoody Fm. (USA)	Griesbachian – Dienerian	15	Hofmann <i>et al.</i> (2013 <i>a</i> )
Köveskál Fm., Arács Fm., Alcsútdoboz Fm. (Balaton Highland, Hungary)	Griesbachian – Dienerian	12	Broglia Loriga <i>et al.</i> (1990)
Lazurnaya Bay Fm. (Russia)	Griesbachian	12	Shigeta <i>et al.</i> (2009)
Bódvaszilas Sandstone Fm. (Hungary)	middle Griesbachian – Smithian	12	Foster <i>et al.</i> (2015)
Luolou Fm. (China)	middle Griesbachian	20	Hautmann <i>et al.</i> (2015)
Andraz Horizon (Werfen Fm., Italy)	middle Griesbachian	3	Foster <i>et al.</i> (2017)
Mazzin Mb. (Werfen Fm., Italy)	early – middle Griesbachian	12	Broglia Loriga <i>et al.</i> (1986, 1990), Hofmann <i>et al.</i> (2015), Foster <i>et</i> <i>al.</i> (2017)
microbialite unit	early Griesbachian	26	Foster <i>et al.</i> (2019)

(Daye Fm., China)			
Tesero Oolite (Werfen Fm., Italy)	late Changhsingian – early Griesbachian	5	Broglia Loriga <i>et al.</i> (1986, 1990), Foster <i>et al.</i> (2017)